

PRKAB2

Purified Mouse Monoclonal Antibody Catalog # AO2604a

Specification

PRKAB2 - Product Information

Application WB, IHC, ICC, E

Primary Accession
Reactivity
Host
Clonality
Monoclonal
Isotype
Calculated MW

Muse
Mouse IgG2b
Mouse IgG2b
Mouse IgG2b

Immunogen

Purified recombinant fragment of human PRKAB2 (AA: 1-120) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

PRKAB2 - Additional Information

Gene ID 5565

Dilution

WB~~ 1/500 - 1/2000 IHC~~1:100~500 ICC~~N/A E~~ 1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PRKAB2 is for research use only and not for use in diagnostic or therapeutic procedures.

PRKAB2 - Protein Information

Name PRKAB2

Function

Non-catalytic subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of



cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Beta non-catalytic subunit acts as a scaffold on which the AMPK complex assembles, via its C-terminus that bridges alpha (PRKAA1 or PRKAA2) and gamma subunits (PRKAG1, PRKAG2 or PRKAG3).

PRKAB2 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

PRKAB2 - Images




Figure 1:Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)






Figure 2:Western blot analysis using PRKAB2 mAb against human PRKAB2 (AA: 1-120) recombinant protein. (Expected MW is 39 kDa)




Figure 3:Western blot analysis using PRKAB2 mAb against HEK293 (1) and PRKAB2 (AA: 1-120)-hlgGFc transfected HEK293 (2) cell lysate.




Figure 4:Flow cytometric analysis of Hela cells using PRKAB2 mouse mAb (green) and negative control (red).

PRKAB2 - References

1.Mol Biol Cell. 2013 Jun;24(11):1801-11, S1-4.2.Circ Res. 2012 Aug 31;111(6):800-14.