

MET10 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10007B

Specification

MET10 Antibody (C-term) - Product Information

IHC-P, WB,E Application **Primary Accession** 086W50 Other Accession NP 076991.3 Human, Mouse Reactivity Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Antigen Region 444-472

MET10 Antibody (C-term) - Additional Information

Gene ID 79066

Other Names

Methyltransferase-like protein 16, 211-, Methyltransferase 10 domain-containing protein, METTL16, METT10D

Target/Specificity

This MET10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 444-472 amino acids from the C-terminal region of human MET10.

Dilution

IHC-P~~1:100 WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MET10 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MET10 Antibody (C-term) - Protein Information

Name METTL16 {ECO:0000303|PubMed:27872311, ECO:0000312|HGNC:HGNC:28484}



Function RNA N6-methyltransferase that methylates adenosine residues at the N(6) position of a subset of RNAs and is involved in S-adenosyl- L-methionine homeostasis by regulating expression of MAT2A transcripts (PubMed: 28525753, PubMed: 30197297, PubMed: 30197299, PubMed:33428944, PubMed:33930289). Able to N6-methylate a subset of mRNAs and U6 small nuclear RNAs (U6 snRNAs) (PubMed: 28525753). In contrast to the METTL3- METTL14 heterodimer, only able to methylate a limited number of RNAs: requires both a 5'UACAGAGAA-3' nonamer sequence and a specific RNA structure (PubMed: 28525753, PubMed: 30197297, PubMed:30197299). Plays a key role in S-adenosyl-L-methionine homeostasis by mediating N6methylation of MAT2A mRNAs, altering splicing of MAT2A transcripts: in presence of S-adenosyl-L-methionine, binds the 3'-UTR region of MAT2A mRNA and specifically N6-methylates the first hairpin of MAT2A mRNA, preventing recognition of their 3'-splice site by U2AF1/U2AF35, thereby inhibiting splicing and protein production of S-adenosylmethionine synthase (PubMed: 28525753, PubMed: 33930289). In S-adenosyl-L- methionine-limiting conditions, binds the 3'-UTR region of MAT2A mRNA but stalls due to the lack of a methyl donor, preventing N6-methylation and promoting expression of MAT2A (PubMed: 28525753). In addition to mRNAs, also able to mediate N6-methylation of U6 small nuclear RNA (U6 snRNA): specifically N6-methylates adenine in position 43 of U6 snRNAs (PubMed: 28525753, PubMed: 29051200, PubMed: 32266935). Also able to bind various IncRNAs, such as 7SK snRNA (7SK RNA) or 7SL RNA (PubMed: 29051200). Specifically binds the 3'-end of the MALAT1 long non-coding RNA (PubMed: 27872311).

Cellular Location Nucleus. Cytoplasm


MET10 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MET10 Antibody (C-term) - Images





Anti-MET10 Antibody (C-term) at 1:2000 dilution + Mouse bladder lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 64 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of AP10007B on paraffin-embedded Human bladder tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.





Immunohistochemical analysis of AP10007B on paraffin-embedded Human brain tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.