

KIAA1680 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10022b**Specification**

KIAA1680 Antibody (C-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	O9C0I3
Other Accession	NP_001138537.1 , NP_997374.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	99510
Antigen Region	780-808

KIAA1680 Antibody (C-term) - Additional Information**Gene ID** 401145**Other Names**

Serine-rich coiled-coil domain-containing protein 1, Coiled-coil serine-rich protein 1, CCSER1, FAM190A, KIAA1680

Target/Specificity

This KIAA1680 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 780-808 amino acids from the C-terminal region of human KIAA1680.

Dilution

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KIAA1680 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

KIAA1680 Antibody (C-term) - Protein Information**Name** CCSER1


Synonyms FAM190A, KIAA1680

KIAA1680 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KIAA1680 Antibody (C-term) - Images



KIAA1680 Antibody (C-term) (Cat. #AP10022b) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the KIAA1680 antibody detected the KIAA1680 protein (arrow).



KIAA1680 antibody (C-term) (Cat. #AP10022b) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the KIAA1680 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

