

**KEAP1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10092b****Specification**

---

**KEAP1 Antibody (C-term) - Product Information**

Application	IHC-P, WB,E
Primary Accession	<a href="#">Q14145</a>
Other Accession	<a href="#">Q684M4</a> , <a href="#">NP_036421.2</a> , <a href="#">NP_987096.1</a>
Reactivity	Human
Predicted	Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	69666
Antigen Region	422-449

**KEAP1 Antibody (C-term) - Additional Information****Gene ID** 9817**Other Names**

Kelch-like ECH-associated protein 1, Cytosolic inhibitor of Nrf2, INrf2, Kelch-like protein 19, KEAP1, INRF2, KIAA0132, KLHL19

**Target/Specificity**

This KEAP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 422-449 amino acids from the C-terminal region of human KEAP1.

**Dilution**

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

KEAP1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**KEAP1 Antibody (C-term) - Protein Information**

**Name** KEAP1 {ECO:0000303|PubMed:14585973, ECO:0000312|HGNC:HGNC:23177}

**Function** Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin ligase complex that regulates the response to oxidative stress by targeting NFE2L2/NRF2 for ubiquitination (PubMed:[14585973](#), PubMed:[15379550](#), PubMed:[15572695](#), PubMed:[15601839](#), PubMed:[15983046](#), PubMed:[37339955](#)). KEAP1 acts as a key sensor of oxidative and electrophilic stress: in normal conditions, the BCR(KEAP1) complex mediates ubiquitination and degradation of NFE2L2/NRF2, a transcription factor regulating expression of many cytoprotective genes (PubMed:[15601839](#), PubMed:[16006525](#)). In response to oxidative stress, different electrophile metabolites trigger non-enzymatic covalent modifications of highly reactive cysteine residues in KEAP1, leading to inactivate the ubiquitin ligase activity of the BCR(KEAP1) complex, promoting NFE2L2/NRF2 nuclear accumulation and expression of phase II detoxifying enzymes (PubMed:[16006525](#), PubMed:[17127771](#), PubMed:[18251510](#), PubMed:[19489739](#), PubMed:[29590092](#)). In response to selective autophagy, KEAP1 is sequestered in inclusion bodies following its interaction with SQSTM1/p62, leading to inactivation of the BCR(KEAP1) complex and activation of NFE2L2/NRF2 (PubMed:[20452972](#)). The BCR(KEAP1) complex also mediates ubiquitination of SQSTM1/p62, increasing SQSTM1/p62 sequestering activity and degradation (PubMed:[28380357](#)). The BCR(KEAP1) complex also targets BPTF and PGAM5 for ubiquitination and degradation by the proteasome (PubMed:[15379550](#), PubMed:[17046835](#)).

#### **Cellular Location**

Cytoplasm. Nucleus. Note=Mainly cytoplasmic (PubMed:15601839). In response to selective autophagy, relocates to inclusion bodies following interaction with SQSTM1/p62 (PubMed:20452972).

#### **Tissue Location**


Broadly expressed, with highest levels in skeletal muscle.


#### **KEAP1 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **KEAP1 Antibody (C-term) - Images**





KEAP1 Antibody (C-term) (Cat. #AP10092b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the KEAP1 antibody detected the KEAP1 protein (arrow).



KEAP1 antibody (C-term) (Cat. #AP10092b) immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the KEAP1 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.