

STIM1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10114b**Specification**

STIM1 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q13586
Other Accession	P70302 , NP_003147.2
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	77423
Antigen Region	509-538

STIM1 Antibody (C-term) - Additional Information**Gene ID** 6786**Other Names**

Stromal interaction molecule 1, STIM1, GOK

Target/Specificity

This STIM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 509-538 amino acids from the C-terminal region of human STIM1.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

STIM1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

STIM1 Antibody (C-term) - Protein Information**Name** STIM1

Synonyms GOK {ECO:0000303|PubMed:9377559}

Function Plays a role in mediating store-operated Ca^{2+} entry (SOCE), a Ca^{2+} influx following depletion of intracellular Ca^{2+} stores (PubMed:[15866891](#), PubMed:[16005298](#), PubMed:[16208375](#), PubMed:[16537481](#), PubMed:[16733527](#), PubMed:[16766533](#), PubMed:[16807233](#), PubMed:[18854159](#), PubMed:[19249086](#), PubMed:[22464749](#), PubMed:[24069340](#), PubMed:[24351972](#), PubMed:[24591628](#), PubMed:[26322679](#), PubMed:[25326555](#), PubMed:[28219928](#)). Acts as a Ca^{2+} sensor in the endoplasmic reticulum via its EF-hand domain. Upon Ca^{2+} depletion, translocates from the endoplasmic reticulum to the plasma membrane where it activates the Ca^{2+} release-activated Ca^{2+} (CRAC) channel subunit ORAI1 (PubMed:[16208375](#), PubMed:[16537481](#)). Involved in enamel formation (PubMed:[24621671](#)). Activated following interaction with STIMATE, leading to promote STIM1 conformational switch (PubMed:[26322679](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Cytoplasm, cytoskeleton. Sarcoplasmic reticulum.

Note=Translocates from the endoplasmic reticulum to the cell membrane in response to a depletion of intracellular calcium and is detected at punctae corresponding to junctions between the endoplasmic reticulum and the cell membrane (PubMed:[19249086](#), PubMed:[16005298](#), PubMed:[16208375](#), PubMed:[18854159](#)) Associated with the microtubule network at the growing distal tip of microtubules (PubMed:[19632184](#)). Colocalizes with ORAI1 at the cell membrane (PubMed:[27185316](#)). Colocalizes preferentially with CASQ1 at endoplasmic reticulum in response to a depletion of intracellular calcium (PubMed:[27185316](#)).

Tissue Location

Ubiquitously expressed in various human primary cells and tumor cell lines.


STIM1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STIM1 Antibody (C-term) - Images





STIM1 Antibody (C-term) (Cat. #AP10114b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the STIM1 antibody detected the STIM1 protein (arrow).



STIM1 Antibody (C-term) (Cat. #AP10114b) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the STIM1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

STIM1 Antibody (C-term) - Background

This gene encodes a type 1 transmembrane protein that mediates Ca^{2+} influx after depletion of intracellular Ca^{2+} stores by gating of store-operated Ca^{2+} influx channels (SOCs). It is one of several genes located in the imprinted gene domain of 11p15.5, an important tumor-suppressor gene region. Alterations in this region have been associated with the Beckwith-Wiedemann syndrome, Wilms tumor, rhabdomyosarcoma, adrenocortical carcinoma, and lung, ovarian, and breast cancer. This gene may play a role in malignancies and disease that involve this region, as well as early hematopoiesis, by mediating attachment to stromal cells. This gene is oriented in a head-to-tail configuration with the ribonucleotide reductase 1 gene (RRM1), with the 3' end of this gene situated 1.6 kb from the 5' end of the RRM1 gene.

STIM1 Antibody (C-term) - References

Byun, M., et al. J. Exp. Med. 207(11):2307-2312(2010)
Park, C.Y., et al. Science 330(6000):101-105(2010)
Walsh, C.M., et al. Biochem. J. 430(3):453-460(2010)

Hawkins, B.J., et al. J. Cell Biol. 190(3):391-405(2010)

Woodward, O.M., et al. PLoS ONE 5 (8), E12305 (2010) :

STIM1 Antibody (C-term) - Citations

- [Suppression of STIM1 inhibits the migration and invasion of human prostate cancer cells and is associated with PI3K/Akt signaling inactivation.](#)