

FOXL2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10166A**Specification**

FOXL2 Antibody (N-term) - Product Information

Application	FC, WB,E
Primary Accession	P58012
Other Accession	NP_075555.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	38772
Antigen Region	3-29

FOXL2 Antibody (N-term) - Additional Information**Gene ID** 668**Other Names**

Forkhead box protein L2, FOXL2

Target/Specificity

This FOXL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 3-29 amino acids from the N-terminal region of human FOXL2.

Dilution

FC~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FOXL2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FOXL2 Antibody (N-term) - Protein Information**Name** FOXL2

Function Transcriptional regulator. Critical factor essential for ovary differentiation and maintenance, and repression of the genetic program for somatic testis determination. Prevents trans- differentiation of ovary to testis through transcriptional repression of the Sertoli cell-promoting gene SOX9 (By similarity). Has apoptotic activity in ovarian cells. Suppresses ESR1-mediated transcription of PTGS2/COX2 stimulated by tamoxifen (By similarity). Is a regulator of CYP19 expression (By similarity). Participates in SMAD3-dependent transcription of FST via the intronic SMAD-binding element (By similarity). Is a transcriptional repressor of STAR. Activates SIRT1 transcription under cellular stress conditions. Activates transcription of OSR2.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00089, ECO:0000269|PubMed:19744555}

Tissue Location


In addition to its expression in the developing eyelid, it is transcribed very early in somatic cells of the developing gonad (before sex determination) and its expression persists in the follicular cells of the adult ovary

FOXL2 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FOXL2 Antibody (N-term) - Images



FOXL2 Antibody (N-term) (Cat. #AP10166a) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the FOXL2 antibody detected the FOXL2 protein (arrow).



FOXL2 Antibody (N-term) (Cat. #AP10166a) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

FOXL2 Antibody (N-term) - Background

This gene encodes a forkhead transcription factor. The protein contains a fork-head DNA-binding domain and may play a role in ovarian development and function. Mutations in this gene are a cause of blepharophimosis syndrome and premature ovarian failure 3.

FOXL2 Antibody (N-term) - References

Nakashima, M., et al. Nat. Genet. 42(9):768-771(2010) Kraoua, L., et al. Clin. Genet. 77(6):601-603(2010) Murphy, B.D. Nat. Med. 16(5):527-529(2010) Corpuz, P.S., et al. Mol. Endocrinol. 24(5):1037-1051(2010) Kim, T., et al. Histopathology 56(3):408-410(2010)

FOXL2 Antibody (N-term) - Citations

- [The Emerging Role of FOXL2 in Regulating the Transcriptional Activation Function of Estrogen Receptor \$\beta\$: An Insight Into Ovarian Folliculogenesis.](#)