

**GABRG2 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10189c**

**Specification**

**GABRG2 Antibody (Center) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">P18507</a>
Other Accession	<a href="#">P18508</a> , <a href="#">P22723</a> , <a href="#">P22300</a> , <a href="#">NP_000807.2</a>
Reactivity	Human, Mouse
Predicted	Bovine, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	198-224

**GABRG2 Antibody (Center) - Additional Information**

**Gene ID 2566**

**Other Names**

Gamma-aminobutyric acid receptor subunit gamma-2, GABA(A) receptor subunit gamma-2,  
GABRG2

**Target/Specificity**

This GABRG2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 198-224 amino acids from the Central region of human GABRG2.

**Dilution**

WB~~1:1000

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GABRG2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**GABRG2 Antibody (Center) - Protein Information**

Name GABRG2 ([HGNC:4087](#))

**Function** Gamma subunit of the heteropentameric ligand-gated chloride channel gated by gamma-aminobutyric acid (GABA), a major inhibitory neurotransmitter in the brain (PubMed:[14993607](#), PubMed:[16412217](#), PubMed:[23909897](#), PubMed:[2538761](#), PubMed:[25489750](#), PubMed:[27864268](#), PubMed:[29950725](#), PubMed:[30602789](#)). GABA-gated chloride channels, also named GABA(A) receptors (GABAAR), consist of five subunits arranged around a central pore and contain GABA active binding site(s) located at the alpha and beta subunit interface(s) (PubMed:[29950725](#), PubMed:[30602789](#)). When activated by GABA, GABAARs selectively allow the flow of chloride anions across the cell membrane down their electrochemical gradient (PubMed:[14993607](#), PubMed:[16412217](#), PubMed:[2538761](#), PubMed:[27864268](#), PubMed:[29950725](#), PubMed:[30602789](#)). Gamma-2/GABRG2-containing GABAARs are found at both synaptic and extrasynaptic sites (By similarity). Chloride influx into the postsynaptic neuron following GABAAR opening decreases the neuron ability to generate a new action potential, thereby reducing nerve transmission (By similarity). GABAARs containing alpha-1 and beta-2 or -3 subunits exhibit synaptogenic activity; the gamma-2 subunit being necessary but not sufficient to induce rapid synaptic contacts formation (PubMed:[23909897](#), PubMed:[25489750](#)). Extrasynaptic gamma-2-containing receptors contribute to the tonic GABAergic inhibition (By similarity). GABAARs function also as histamine receptor where histamine binds at the interface of two neighboring beta subunits and potentiates GABA response in a gamma-2 subunit-controlled manner (By similarity).

#### Cellular Location


Postsynaptic cell membrane; Multi-pass membrane protein {ECO:0000269|PubMed:30602789, ECO:0007744|PDB:6I53}. Cell membrane; Multi-pass membrane protein {ECO:0000269|PubMed:30602789, ECO:0007744|PDB:6I53} Cell projection, dendrite {ECO:0000250|UniProtKB:P22723}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:P18508}

#### GABRG2 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### GABRG2 Antibody (Center) - Images



GABRG2 Antibody (Center) (Cat. #AP10189c) western blot analysis in mouse testis tissue lysates (15ug/lane). This demonstrates the GABRG2 antibody detected GABRG2 protein (arrow).



Anti-GABRG2 Antibody (Center) at 1:1000 dilution + human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 54 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



GABRG2 antibody (Center) (Cat. #AP10189c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the GABRG2 antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **GABRG2 Antibody (Center) - Background**

This gene encodes a gamma-aminobutyric acid (GABA) receptor. GABA is the major inhibitory neurotransmitter in the mammalian brain, where it acts at GABA-A receptors, which are ligand-gated chloride channels. GABA-A receptors are pentameric, consisting of proteins from several subunit classes: alpha, beta, gamma, delta and rho. Mutations in this gene have been associated with epilepsy and febrile seizures. Multiple transcript variants encoding different isoforms have been identified for this gene.

#### **GABRG2 Antibody (Center) - References**

Green, E.K., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 153B (7), 1347-1349 (2010) :

Jansen, L.A., et al. Epilepsia 51(8):1456-1467(2010)  
Pinheiro, A.P., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 153B (5), 1070-1080 (2010) :  
Shi, X., et al. J. Hum. Genet. 55(6):375-378(2010)  
Kumari, R., et al. Seizure 19(4):237-241(2010)