

**GCC2 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10197b****Specification**

---

**GCC2 Antibody (C-term) - Product Information**

Application	IHC-P, WB,E
Primary Accession	<a href="#">Q8IWJ2</a>
Other Accession	<a href="#">O14715</a> , <a href="#">Q99666</a> , <a href="#">D3ZZL9</a> , <a href="#">Q8CHG3</a> , <a href="#">NP_852118.1</a>
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	195910
Antigen Region	1581-1609

**GCC2 Antibody (C-term) - Additional Information****Gene ID** 9648**Other Names**

GRIP and coiled-coil domain-containing protein 2, 185 kDa Golgi coiled-coil protein, GCC185, CLL-associated antigen KW-11, CTCL tumor antigen se1-1, Ran-binding protein 2-like 4, RanBP2L4, Renal carcinoma antigen NY-REN-53, GCC2, KIAA0336, RANBP2L4

**Target/Specificity**

This GCC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1581-1609 amino acids from the C-terminal region of human GCC2.

**Dilution**

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GCC2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**GCC2 Antibody (C-term) - Protein Information**

**Name** GCC2

**Synonyms** KIAA0336, RANBP2L4

**Function** Golgin which probably tethers transport vesicles to the trans-Golgi network (TGN) and regulates vesicular transport between the endosomes and the Golgi. As a RAB9A effector it is involved in recycling of the mannose 6-phosphate receptor from the late endosomes to the TGN. May also play a role in transport between the recycling endosomes and the Golgi. Required for maintenance of the Golgi structure, it is involved in the biogenesis of noncentrosomal, Golgi-associated microtubules through recruitment of CLASP1 and CLASP2.

**Cellular Location**

Cytoplasm. Golgi apparatus, trans-Golgi network membrane; Peripheral membrane protein

**Tissue Location**


Ubiquitous..

**GCC2 Antibody (C-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**GCC2 Antibody (C-term) - Images**



GCC2 Antibody (C-term) (Cat. #AP10197b) western blot analysis in mouse stomach tissue lysates (15ug/lane). This demonstrates the GCC2 antibody detected GCC2 protein (arrow).



GCC2 antibody (C-term) (Cat. #AP10197b) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the GCC2 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **GCC2 Antibody (C-term) - Background**

The protein encoded by this gene is a peripheral membrane protein localized to the trans-Golgi network. It is sensitive to brefeldin A. This encoded protein contains a GRIP domain which is thought to be used in targeting. Alternative splicing results in multiple transcript variants.

#### **GCC2 Antibody (C-term) - References**

Houghton, F.J., et al. Cell 138(4):787-794(2009)  
Hayes, G.L., et al. Mol. Biol. Cell 20(1):209-217(2009)  
Burguete, A.S., et al. Cell 132(2):286-298(2008)  
Efimov, A., et al. Dev. Cell 12(6):917-930(2007)  
Derby, M.C., et al. Traffic 8(6):758-773(2007)