

HIF1AN Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP1029b

Specification

HIF1AN Antibody (C-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region WB,E <u>O9NWT6</u> <u>O8BLR9</u>, <u>P59723</u> Human, Mouse Zebrafish Rabbit Polyclonal Rabbit IgG 40285 286-316

HIF1AN Antibody (C-term) - Additional Information

Gene ID 55662

Other Names Hypoxia-inducible factor 1-alpha inhibitor, 11411n4, Factor inhibiting HIF-1, FIH-1, Hypoxia-inducible factor asparagine hydroxylase, HIF1AN, FIH1

Target/Specificity

This HIF1AN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 286-316 amino acids from the C-terminal region of human HIF1AN.

Dilution

WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HIF1AN Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HIF1AN Antibody (C-term) - Protein Information

Name HIF1AN



Synonyms FIH1

Function Hydroxylates HIF-1 alpha at 'Asn-803' in the C-terminal transactivation domain (CAD). Functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents interaction of HIF-1 with transcriptional coactivators including Cbp/p300-interacting transactivator. Involved in transcriptional repression through interaction with HIF1A, VHL and histone deacetylases. Hydroxylates specific Asn residues within ankyrin repeat domains (ARD) of NFKB1, NFKBIA, NOTCH1, ASB4, PPP1R12A and several other ARD-containing proteins. Also hydroxylates Asp and His residues within ARDs of ANK1 and TNKS2, respectively. Negatively regulates NOTCH1 activity, accelerating myogenic differentiation. Positively regulates ASB4 activity, promoting vascular differentiation.

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, perinuclear region. Note=Mainly cytoplasmic localization, but interaction with NOTCH1 results in nuclear localization and interaction with ABPA3 results in perinuclear localization in macrophages

HIF1AN Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

HIF1AN Antibody (C-term) - Images

A375	m.brain
130	150
72 55 - 4	75
36	37
20	25
17	20
11	15

Western blot analysis of anti-HIF1AN (C-term) Pab in A375 and mouse brain cell line lysate. HIF1AN (C-term)(arrow) was detected using the purified Pab.

HIF1AN Antibody (C-term) - Background

FIH1Encoded protein (factor inhibiting HIF-1)is a co-repressor that interacts with hypoxia-inducible factor 1(HIF-1) alpha and the von Hippel-Lindau tumor suppressor protein to mediate repression of HIF-1 transcriptional activity.



HIF1AN Antibody (C-term) - Citations

- Dickkopf-related protein 3 alters aerobic glycolysis in pancreatic cancer BxPC-3 cells, promoting CD4 T-cell activation and function
- miR-148a-3p Mediates Notch Signaling to Promote the Differentiation and M1 Activation of Macrophages.