

MBD1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP1036B

Specification

MBD1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	<u>Q9UIS9</u>
Reactivity	Human
Host	Rabbit
Clonality	Polycional
Isotype	Rabbit IgG
Antigen Region	376-405

MBD1 Antibody (C-term) - Additional Information

Gene ID 4152

Other Names Methyl-CpG-binding domain protein 1, CXXC-type zinc finger protein 3, Methyl-CpG-binding protein MBD1, Protein containing methyl-CpG-binding domain 1, MBD1, CXXC3, PCM1

Target/Specificity

This MBD1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 376-405 amino acids from the C-terminal region of human MBD1.

Dilution WB~~1:1000

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MBD1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MBD1 Antibody (C-term) - Protein Information

Name MBD1 (HGNC:6916)

Synonyms CXXC3, PCM1

Function Transcriptional repressor that binds CpG islands in promoters where the DNA is



methylated at position 5 of cytosine within CpG dinucleotides. Binding is abolished by the presence of 7-mG that is produced by DNA damage by methylmethanesulfonate (MMS). Acts as transcriptional repressor and plays a role in gene silencing by recruiting ATF7IP, which in turn recruits factors such as the histone methyltransferase SETDB1. Probably forms a complex with SETDB1 and ATF7IP that represses transcription and couples DNA methylation and histone 'Lys-9' trimethylation. Isoform 1 and isoform 2 can also repress transcription from unmethylated promoters.

Cellular Location

Nucleus. Nucleus matrix. Nucleus speckle Chromosome Note=Nuclear, in a punctate pattern (PubMed:12711603). Associated with euchromatic regions of the chromosomes, with pericentromeric regions on chromosome 1 and with telomeric regions from several chromosomes (PubMed:10454587, PubMed:10648624).

Tissue Location Widely expressed..

MBD1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

MBD1 Antibody (C-term) - Images



Western blot analysis of anti-MBD1 Pab (Cat. #AP1036b) in HeLa cell lysate. MBD1 (Arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

MBD1 Antibody (C-term) - Background

DNA methylation, or the addition of methyl groups to cytosine bases in the dinucleotide CpG, is



imperative to proper development and regulates gene expression. The methylation pattern involves the enzymatic processes of methylation and demethylation. The demethylation enzyme was recently found

to be a mammalian protein, which exhibits demethylase activity associated to a

methyl-CpG-binding domain (MBD) (1). The enzyme is able to revert methylated cytosine bases to cytosines within the particular dinucleotide sequence mdCpdG by catalyzing the cleaving of the methyl group as methanol.

MeCP2 and MBD1 (PCM1) are first found to repress transcription by binding specifically to methylated DNA (2). MBD2 and MBD4 (also known as MED1) were later found to colocalize with foci of heavily methylated satellite DNA and

believed to mediate the biological functions of the methylation signal. Surprisingly, MBD3 does not bind methylated DNA both in vivo and in vitro. MBD1, MBD2, MBD3, and MBD4 are found to be expressed in somatic tissues, but the expression of MBD1 and MBD2 is reduced or absent in embryonic stem

cells, which are known to be deficient in MeCP1 activity. MBD4 have homology to bacterial base excision repair DNA N-glycosylases/lyases (3). In some microsatellite unstable tumors MBD4 is mutated at an exonic polynucleotide tract (4).

MBD1 Antibody (C-term) - References

Bhattacharya SK, Ramchandani S, Cervoni N, Szyf. M. Nature, 397 (6720):579-583 1999. Hendrich B and Bird A. Mol Cell Biol, 18: 6538-6547(1998).

Petronzelli F, Riccio A, Markham GD, Seeholzer SH, Stoerker J, Genuardi M, Yeung AT, Matsumoto Y, Bellacosa A. J Biol Chem 275 (42): 32422-32429 (2000).

Bader S, Walker M, Harrison D. Br J Cancer 83(12): 1646-1649 (2000).

MBD1 Antibody (C-term) - Citations

- Specific binding of the methyl binding domain protein 2 at the BRCA1-NBR2 locus.
- The intracellular domain of teneurin-1 interacts with MBD1 and CAP/ponsin resulting in subcellular codistribution and translocation to the nuclear matrix.