

KIAA1644 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10426a**Specification**

KIAA1644 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q3SXP7
Other Accession	Q0VBP7 , NP_001092764.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	22573
Antigen Region	18-46

KIAA1644 Antibody (N-term) - Additional Information**Gene ID** 85352**Other Names**

Uncharacterized protein KIAA1644, KIAA1644

Target/Specificity

This KIAA1644 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 18-46 amino acids from the N-terminal region of human KIAA1644.

Dilution

WB~~1:1000

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KIAA1644 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

KIAA1644 Antibody (N-term) - Protein Information**Name** SHISAL1 ([HGNC:29335](#))


Synonyms KIAA1644**Cellular Location**

Membrane; Single-pass type I membrane protein


KIAA1644 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KIAA1644 Antibody (N-term) - Images

KIAA1644 Antibody (N-term) (Cat. #AP10426a) western blot analysis in mouse bladder tissue lysates (35ug/lane). This demonstrates the KIAA1644 antibody detected the KIAA1644 protein (arrow).



KIAA1644 antibody (N-term) (Cat. #AP10426a) immunohistochemistry analysis in formalin fixed and paraffin embedded human uterus tissue followed by peroxidase conjugation of the secondary

antibody and DAB staining. This data demonstrates the use of the KIAA1644 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

KIAA1644 Antibody (N-term) - References

Hirosawa, M., et al. DNA Res. 8(1):1-9(2001)