

CHIC2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10473b**Specification**

CHIC2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9UKJ5
Other Accession	O9D9G3 , NP_036242.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	19254
Antigen Region	138-165

CHIC2 Antibody (C-term) - Additional Information**Gene ID** 26511**Other Names**

Cysteine-rich hydrophobic domain-containing protein 2, BrX-like translocated in leukemia, CHIC2, BTL

Target/Specificity

This CHIC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 138-165 amino acids from the C-terminal region of human CHIC2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CHIC2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CHIC2 Antibody (C-term) - Protein Information**Name** CHIC2


Synonyms BTL**Cellular Location**

Cell membrane. Cytoplasmic vesicle. Note=Also present at a Golgi-like vesicular compartment and at scattered vesicles

CHIC2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CHIC2 Antibody (C-term) - Images

CHIC2 Antibody (C-term) (Cat. #AP10473b) western blot analysis in mouse bladder tissue lysates (35ug/lane). This demonstrates the CHIC2 antibody detected the CHIC2 protein (arrow).

CHIC2 Antibody (C-term) - Background

CHIC2 is a member of the CHIC family of proteins. This protein contains a cysteine-rich hydrophobic (CHIC) motif, and is localized to vesicular structures and the plasma membrane.

CHIC2 Antibody (C-term) - References

Fink, S.R., et al. Leuk. Res. 33(6):843-846(2009)
Venkatesan, K., et al. Nat. Methods 6(1):83-90(2009)
Pardanani, A., et al. Blood 102(9):3093-3096(2003)
Cools, J., et al. FEBS Lett. 492(3):204-209(2001)
Cools, J., et al. FEBS Lett. 492(3):204-209(2001)