

INPP5B Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10490b

Specification

INPP5B Antibody (C-term) - Product Information

Application WB, IHC-P, FC,E **Primary Accession** P32019 Other Accession NP 005531.2 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 112852 Antigen Region 950-979

INPP5B Antibody (C-term) - Additional Information

Gene ID 3633

Other Names

Type II inositol 1, 5-trisphosphate 5-phosphatase, 75 kDa inositol polyphosphate-5-phosphatase, Phosphoinositide 5-phosphatase, 5PTase, INPP5B, OCRL2

Target/Specificity

This INPP5B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 950-979 amino acids from the C-terminal region of human INPP5B.

Dilution

WB~~1:1000 IHC-P~~1:50~100 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

INPP5B Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

INPP5B Antibody (C-term) - Protein Information

Name INPP5B





Synonyms OCRL2

Function Hydrolyzes phosphatidylinositol 4,5-bisphosphate (PtIns(4,5)P2) and the signaling molecule phosphatidylinositol 1,4,5- trisphosphate (PtIns(1,4,5)P3), and thereby modulates cellular signaling events.

Cellular Location

Cytoplasm, cytosol. Endoplasmic reticulum-Golgi intermediate compartment. Early endosome membrane. Membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:Q8K337}. Golgi apparatus

Tissue Location


Platelets.

INPP5B Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

INPP5B Antibody (C-term) - Images




Anti-INPP5B Antibody (C-term) at 1:1000 dilution + Daudi whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 113 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





INPP5B antibody (C-term) (Cat. #AP10490b) immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the INPP5B antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.




INPP5B Antibody (C-term) (Cat. #AP10490b) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

INPP5B Antibody (C-term) - Background

Cellular calcium signaling is controlled by the production of inositol phosphates (IPs) by phospholipase C in response to extracellular signals. The IP signaling molecules are inactivated by a family of inositol polyphosphate-5-phosphatases (5-phosphatases). INPP5B encodes the type II 5-phosphatase. The protein is localized to the cytosol and mitochondria, and associates with membranes through an isoprenyl modification near the C-terminus. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been determined. [provided by RefSeq].

INPP5B Antibody (C-term) - References

Coon, B.G., et al. Hum. Mol. Genet. 18(23):4478-4491(2009) Mao, Y., et al. EMBO J. 28(13):1831-1842(2009) Williams, C., et al. J. Cell. Sci. 120 (PT 22), 3941-3951 (2007) :





Speed, C.J., et al. Eur. J. Biochem. 234(1):216-224(1995) Janne, P.A., et al. Genomics 28(2):280-285(1995)