

TPSD1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10500b**Specification**

TPSD1 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9BZJ3
Other Accession	NP_036349.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	26584
Antigen Region	164-193

TPSD1 Antibody (C-term) - Additional Information**Gene ID** 23430**Other Names**

Tryptase delta, Delta-tryptase, HmMCP-3-like tryptase III, Mast cell mMCP-7-like, Tryptase-3, TPSD1

Target/Specificity

This TPSD1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 164-193 amino acids from the C-terminal region of human TPSD1.

Dilution

WB~~1:1000

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TPSD1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TPSD1 Antibody (C-term) - Protein Information**Name** TPSD1

Function Trypsase is the major neutral protease present in mast cells and is secreted upon the coupled activation-degranulation response of this cell type.

Cellular Location

Secreted. Note=Released from the secretory granules upon mast cell activation.

Tissue Location


Expressed in colon, lung, heart and synovial tissue. May be specific to mast cells.

TPSD1 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TPSD1 Antibody (C-term) - Images



TPSD1 Antibody (C-term) (Cat. #AP10500b) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the TPSD1 antibody detected the TPSD1 protein (arrow).



TPSD1 antibody (C-term) (Cat. #AP10500b) immunohistochemistry analysis in formalin fixed and paraffin embedded human colon carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TPSD1 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

TPSD1 Antibody (C-term) - Background

Tryptases comprise a family of trypsin-like serine proteases, the peptidase family S1. Tryptases are enzymatically active only as heparin-stabilized tetramers, and they are resistant to all known endogenous proteinase inhibitors. Several tryptase genes are clustered on chromosome 16p13.3. These genes are characterized by several distinct features. They have a highly conserved 3' UTR and contain tandem repeat sequences at the 5' flank and 3' UTR which are thought to play a role in regulation of the mRNA stability. Although this gene may be an exception, most of the tryptase genes have an intron immediately upstream of the initiator Met codon, which separates the site of transcription initiation from protein coding sequence. This feature is characteristic of tryptases but is unusual in other genes. Tryptases have been implicated as mediators in the pathogenesis of asthma and other allergic and inflammatory disorders. TPSD1 was once considered to be a pseudogene, although it is now believed to be a functional gene that encodes a protein.

TPSD1 Antibody (C-term) - References

Jackson, N.E., et al. J. Biol. Chem. 283(49):34178-34187(2008)
Caughey, G.H. J. Allergy Clin. Immunol. 117(6):1411-1414(2006)
Wang, H.W., et al. J. Immunol. 169(9):5145-5152(2002)
Caughey, G.H. Mol. Immunol. 38 (16-18), 1353-1357 (2002) :
Soto, D., et al. Clin. Exp. Allergy 32(7):1000-1006(2002)