

**ZNF98 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10523b**

### Specification

#### ZNF98 Antibody (C-term) - Product Information

|                   |                        |
|-------------------|------------------------|
| Application       | FC, IHC-P, WB,E        |
| Primary Accession | <a href="#">A6NK75</a> |
| Other Accession   | <a href="#">Q9P255</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Isotype           | Rabbit IgG             |
| Calculated MW     | 65799                  |
| Antigen Region    | 520-549                |

#### ZNF98 Antibody (C-term) - Additional Information

##### Gene ID 148198

##### Other Names

Zinc finger protein 98, Zinc finger protein 739, Zinc finger protein F7175, ZNF98, ZNF739

##### Target/Specificity

This ZNF98 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 520-549 amino acids from the C-terminal region of human ZNF98.

##### Dilution

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

##### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

ZNF98 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### ZNF98 Antibody (C-term) - Protein Information

##### Name ZNF98


**Synonyms** ZNF739**Function** May be involved in transcriptional regulation.**Cellular Location**

Nucleus.


**ZNF98 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ZNF98 Antibody (C-term) - Images**

ZNF98 Antibody (C-term) (Cat. #AP10523b) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the ZNF98 antibody detected the ZNF98 protein (arrow).



ZNF98 antibody (C-term) (Cat. #AP10523b) immunohistochemistry analysis in formalin fixed and paraffin embedded human lymphnode followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ZNF98 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



ZNF98 Antibody (C-term) (Cat. #AP10523b) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **ZNF98 Antibody (C-term) - Background**

May be involved in transcriptional regulation.