

ZNF98 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10523b**Specification**

ZNF98 Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	A6NK75
Other Accession	Q9P255
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	65799
Antigen Region	520-549

ZNF98 Antibody (C-term) - Additional Information**Gene ID** 148198**Other Names**

Zinc finger protein 98, Zinc finger protein 739, Zinc finger protein F7175, ZNF98, ZNF739

Target/Specificity

This ZNF98 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 520-549 amino acids from the C-terminal region of human ZNF98.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions


ZNF98 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF98 Antibody (C-term) - Protein Information**Name** ZNF98


Synonyms ZNF739**Function** May be involved in transcriptional regulation.**Cellular Location**
Nucleus.**ZNF98 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF98 Antibody (C-term) - Images

ZNF98 Antibody (C-term) (Cat. #AP10523b) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the ZNF98 antibody detected the ZNF98 protein (arrow).



ZNF98 antibody (C-term) (Cat. #AP10523b) immunohistochemistry analysis in formalin fixed and paraffin embedded human lymphnode followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ZNF98 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



ZNF98 Antibody (C-term) (Cat. #AP10523b) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ZNF98 Antibody (C-term) - Background

May be involved in transcriptional regulation.