

ZNF667 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10552c

Specification

ZNF667 Antibody (Center) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype	FC, IHC-P, WB,E <u>O5HYK9</u> <u>NP_071386.3</u> Human Rabbit Polyclonal Rabbit IgG
	2
Calculated MW	70161
Antigen Region	303-331

ZNF667 Antibody (Center) - Additional Information

Gene ID 63934

Other Names Zinc finger protein 667, ZNF667

Target/Specificity This ZNF667 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 303-331 amino acids from the Central region of human ZNF667.

Dilution FC~~1:10~50 IHC-P~~1:50~100 WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZNF667 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF667 Antibody (Center) - Protein Information

Name ZNF667



Function May be involved in transcriptional regulation.


Cellular Location Nucleus.

ZNF667 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

ZNF667 Antibody (Center) - Images




ZNF667 Antibody (Center) (Cat. #AP10552c) western blot analysis in CEM cell line lysates (35ug/lane).This demonstrates the ZNF667 antibody detected the ZNF667 protein (arrow).



ZNF667 Antibody (Center) (Cat. #AP10552c) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ZNF667 Antibody (Center) for



immunohistochemistry. Clinical relevance has not been evaluated.



ZNF667 Antibody (Center) (Cat. #AP10552c) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ZNF667 Antibody (Center) - Background

May be involved in transcriptional regulation (By similarity).