

ZBTB10 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10573b**Specification**

ZBTB10 Antibody (C-term) - Product Information

Application	FC, WB,E
Primary Accession	O96DT7
Other Accession	NP_076418.3 , NP_001099009.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	94894
Antigen Region	553-582

ZBTB10 Antibody (C-term) - Additional Information**Gene ID** 65986**Other Names**

Zinc finger and BTB domain-containing protein 10, Zinc finger protein RIN ZF, ZBTB10, RINZF, RINZFC

Target/Specificity

This ZBTB10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 553-582 amino acids from the C-terminal region of human ZBTB10.

Dilution

FC~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZBTB10 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZBTB10 Antibody (C-term) - Protein Information**Name** ZBTB10

Synonyms RINZF, RINZFC

Function May be involved in transcriptional regulation.


Cellular Location
Nucleus.

ZBTB10 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZBTB10 Antibody (C-term) - Images



ZBTB10 Antibody (C-term) (Cat. #AP10573b) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the ZBTB10 antibody detected the ZBTB10 protein (arrow).



ZBTB10 Antibody (C-term) (Cat. #AP10573b) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ZBTB10 Antibody (C-term) - Background

May be involved in transcriptional regulation.

ZBTB10 Antibody (C-term) - References

Li, X., et al. Endocrinology 151(6):2462-2473(2010)