

GOLGA2L1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10576b

Specification

GOLGA2L1 Antibody (C-term) - Product Information

Application	FC, WB,E
Primary Accession	O9HBQ8
Other Accession	NP_060070.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	15482
Antigen Region	83-112

GOLGA2L1 Antibody (C-term) - Additional Information

Other Names

Putative golgin subfamily A member 2B, Golgin subfamily A member 2-like protein 1, GOLGA2P5, GOLGA2B, GOLGA2L1

Target/Specificity

This GOLGA2L1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 83-112 amino acids from the C-terminal region of human GOLGA2L1.

Dilution

FC~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GOLGA2L1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GOLGA2L1 Antibody (C-term) - Protein Information

Name GOLGA2P5


Synonyms GOLGA2B, GOLGA2L1

GOLGA2L1 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GOLGA2L1 Antibody (C-term) - Images



GOLGA2L1 Antibody (C-term) (Cat. #AP10576b) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the GOLGA2L1 antibody detected the GOLGA2L1 protein (arrow).



GOLGA2L1 Antibody (C-term) (Cat. #AP10576b) flow cytometric analysis of A549 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

