

**OR51I1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10638b**

### Specification

#### OR51I1 Antibody (C-term) - Product Information

|                   |                                |
|-------------------|--------------------------------|
| Application       | WB,E                           |
| Primary Accession | <a href="#">Q9H343</a>         |
| Other Accession   | <a href="#">NP_001005288.1</a> |
| Reactivity        | Human, Mouse                   |
| Host              | Rabbit                         |
| Clonality         | Polyclonal                     |
| Isotype           | Rabbit IgG                     |
| Calculated MW     | 35484                          |
| Antigen Region    | 284-313                        |

#### OR51I1 Antibody (C-term) - Additional Information

**Gene ID** 390063

#### Other Names

Olfactory receptor 51I1, Odorant receptor HOR5'beta11, Olfactory receptor OR11-39, OR51I1

#### Target/Specificity

This OR51I1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 284-313 amino acids from the C-terminal region of human OR51I1.

#### Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

OR51I1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### OR51I1 Antibody (C-term) - Protein Information

**Name** OR51I1

**Function** Odorant receptor.

## Cellular Location


Cell membrane; Multi-pass membrane protein.

## OR51I1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## OR51I1 Antibody (C-term) - Images



OR51I1 Antibody (C-term) (Cat. #AP10638b) western blot analysis in mouse cerebellum tissue lysates (35ug/lane). This demonstrates the OR51I1 antibody detected the OR51I1 protein (arrow).

## OR51I1 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

## OR51I1 Antibody (C-term) - References

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)  
Bulger, M., et al. Proc. Natl. Acad. Sci. U.S.A. 97(26):14560-14565(2000)  
Bulger, M., et al. Proc. Natl. Acad. Sci. U.S.A. 96(9):5129-5134(1999)