

### WDR76 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10651b

### **Specification**

### WDR76 Antibody (C-term) - Product Information

Application WB,E
Primary Accession Q9H967

Other Accession <u>NP\_001161413.1</u>, <u>NP\_079184.2</u>

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 69769
Antigen Region 455-484

### WDR76 Antibody (C-term) - Additional Information

#### **Gene ID** 79968

#### **Other Names**

WD repeat-containing protein 76, WDR76

#### Target/Specificity

This WDR76 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 455-484 amino acids from the C-terminal region of human WDR76.

## **Dilution**

WB~~1:1000

### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

# **Precautions**

WDR76 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### WDR76 Antibody (C-term) - Protein Information

## Name WDR76

**Function** Specifically binds 5-hydroxymethylcytosine (5hmC), suggesting that it acts as a specific reader of 5hmC.




## WDR76 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## WDR76 Antibody (C-term) - Images



WDR76 Antibody (C-term) (Cat. #AP10651b) western blot analysis in U251 cell line lysates (35ug/lane). This demonstrates the WDR76 antibody detected the WDR76 protein (arrow).



WDR76 Antibody (C-term) (Cat. #AP10651b) western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane). This demonstrates the WDR76 antibody detected the WDR76 protein (arrow).