

## CHRDL1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10667a

## **Specification**

## CHRDL1 Antibody (N-term) - Product Information

**Application** IHC-P, WB,E **Primary Accession** O9BU40 Other Accession NP 660277 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 52026 Antigen Region 2-30

## CHRDL1 Antibody (N-term) - Additional Information

#### **Gene ID 91851**

#### **Other Names**

Chordin-like protein 1, Neuralin-1, Neurogenesin-1, Ventroptin, CHRDL1, NRLN1

#### Target/Specificity

This CHRDL1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 2-30 amino acids from the N-terminal region of human CHRDL1.

# **Dilution**

IHC-P~~1:50~100 WB~~1:1000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

CHRDL1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# CHRDL1 Antibody (N-term) - Protein Information

#### Name CHRDL1



## **Synonyms NRLN1**

**Function** Antagonizes the function of BMP4 by binding to it and preventing its interaction with receptors. Alters the fate commitment of neural stem cells from gliogenesis to neurogenesis. Contributes to neuronal differentiation of neural stem cells in the brain by preventing the adoption of a glial fate. May play a crucial role in dorsoventral axis formation. May play a role in embryonic bone formation (By similarity). May also play an important role in regulating retinal angiogenesis through modulation of BMP4 actions in endothelial cells. Plays a role during anterior segment eye development.

# **Cellular Location** Secreted.

#### **Tissue Location**


Expressed in the developing cornea and in the eye anterior segment in addition to the retina. Differentially expressed in the fetal brain. There is high expression in cerebellum and neocortex Expressed in retinal pericytes.

## CHRDL1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## CHRDL1 Antibody (N-term) - Images



CHRDL1 Antibody (N-term) (Cat. #AP10667a) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the CHRDL1 antibody detected the CHRDL1 protein (arrow).





CHRDL1 antibody (N-term) (Cat. #AP10667a) immunohistochemistry analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CHRDL1 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

# CHRDL1 Antibody (N-term) - Background

CHRDL1 is an antagonist of bone morphogenetic protein 4. The encoded protein may play a role in topographic retinotectal projection and in the regulation of retinal angiogenesis in response to hypoxia. Alternatively spliced transcript variants encoding different isoforms have been described.

## CHRDL1 Antibody (N-term) - References

Fernandes, H., et al. Cells Tissues Organs (Print) 191(6):443-452(2010) Larman, B.W., et al. J. Am. Soc. Nephrol. 20(5):1020-1031(2009) Kane, R., et al. Mol. Vis. 14, 1138-1148 (2008) : Kosinski, C., et al. Proc. Natl. Acad. Sci. U.S.A. 104(39):15418-15423(2007) Gazzerro, E., et al. Rev Endocr Metab Disord 7 (1-2), 51-65 (2006) :