

**ZN572 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP10776b****Specification**

---

**ZN572 Antibody (C-term) - Product Information**

Application	WB, FC,E
Primary Accession	<a href="#">Q7Z3I7</a>
Other Accession	<a href="#">NP_689625.2</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	61238
Antigen Region	471-499

**ZN572 Antibody (C-term) - Additional Information****Gene ID** 137209**Other Names**

Zinc finger protein 572, ZNF572

**Target/Specificity**

This ZN572 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 471-499 amino acids from the C-terminal region of human ZN572.

**Dilution**

WB~~1:1000

FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**


ZN572 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**ZN572 Antibody (C-term) - Protein Information****Name** ZNF572**Function** May be involved in transcriptional regulation.


**Cellular Location**  
Nucleus.**ZN572 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ZN572 Antibody (C-term) - Images**

ZN572 Antibody (C-term) (Cat. #AP10776b) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the ZN572 antibody detected the ZN572 protein (arrow).



ZN572 Antibody (C-term) (Cat. #AP10776b) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

**ZN572 Antibody (C-term) - Background**

May be involved in transcriptional regulation (By similarity).

**ZN572 Antibody (C-term) - References**

Simpson, J.C., et al. EMBO Rep. 1(3):287-292(2000)