

**KAT5 / Tip60/HTATIP Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP1081b**

**Specification**

---

**KAT5 / Tip60/HTATIP Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O92993</a>
Other Accession	<a href="#">O99MK2</a> , <a href="#">O8CHK4</a>
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	58582
Antigen Region	467-498

**KAT5 / Tip60/HTATIP Antibody (C-term) - Additional Information**

**Gene ID** 10524

**Other Names**

Histone acetyltransferase KAT5, 60 kDa Tat-interactive protein, Tip60, Histone acetyltransferase HTATIP, HIV-1 Tat interactive protein, Lysine acetyltransferase 5, cPLA(2)-interacting protein, KAT5, HTATIP, TIP60

**Target/Specificity**

This KAT5 / Tip60/HTATIP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 467-498 amino acids from the C-terminal region of human KAT5 / Tip60/HTATIP.

**Dilution**

WB~~1:1000  
E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

KAT5 / Tip60/HTATIP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**KAT5 / Tip60/HTATIP Antibody (C-term) - Protein Information**

**Name** KAT5 {ECO:0000303|PubMed:32817552, ECO:0000312|HGNC:HGNC:5275}

**Function** Catalytic subunit of the NuA4 histone acetyltransferase complex, a multiprotein complex involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H2A and H4 (PubMed:[12776177](#), PubMed:[14966270](#), PubMed:[15042092](#), PubMed:[15121871](#), PubMed:[15310756](#), PubMed:[16387653](#), PubMed:[19909775](#), PubMed:[25865756](#), PubMed:[27153538](#), PubMed:[29174981](#), PubMed:[29335245](#), PubMed:[32822602](#), PubMed:[33076429](#)). Histone acetylation alters nucleosome-DNA interactions and promotes interaction of the modified histones with other proteins which positively regulate transcription (PubMed:[12776177](#), PubMed:[14966270](#), PubMed:[15042092](#), PubMed:[15121871](#), PubMed:[15310756](#)). The NuA4 histone acetyltransferase complex is required for the activation of transcriptional programs associated with proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair (PubMed:[17709392](#), PubMed:[19783983](#), PubMed:[32832608](#)). The NuA4 complex plays a direct role in repair of DNA double-strand breaks (DSBs) by promoting homologous recombination (HR): the complex inhibits TP53BP1 binding to chromatin via MBTD1, which recognizes and binds histone H4 trimethylated at 'Lys-20' (H4K20me), and KAT5 that catalyzes acetylation of 'Lys-15' of histone H2A (H2AK15ac), thereby blocking the ubiquitination mark required for TP53BP1 localization at DNA breaks (PubMed:[27153538](#), PubMed:[32832608](#)). Also involved in DSB repair by mediating acetylation of 'Lys-5' of histone H2AX (H2AXK5ac), promoting NBN/NBS1 assembly at the sites of DNA damage (PubMed:[17709392](#), PubMed:[26438602](#)). The NuA4 complex plays a key role in hematopoietic stem cell maintenance and is required to maintain acetylated H2A.Z/H2AZ1 at MYC target genes (By similarity). The NuA4 complex is also required for spermatid development by promoting acetylation of histones: histone hyperacetylation is required for histone replacement during the transition from round to elongating spermatids (By similarity). Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AZ1 from the nucleosome (PubMed:[24463511](#)). Also acetylates non-histone proteins, such as BMAL1, ATM, AURKB, CHKA, CGAS, ERCC4/XPF, LPIN1, TP53/p53, NDC80/HEC1, NR1D2, RAN, SOX4, FOXP3, SQSTM1, ULK1 and RUBCNL/Pacer (PubMed:[16141325](#), PubMed:[17189187](#), PubMed:[17360565](#), PubMed:[17996965](#), PubMed:[24835996](#), PubMed:[26829474](#), PubMed:[29040603](#), PubMed:[30409912](#), PubMed:[30704899](#), PubMed:[31857589](#), PubMed:[32034146](#), PubMed:[32817552](#), PubMed:[34077757](#)). Directly acetylates and activates ATM (PubMed:[16141325](#)). Promotes nucleotide excision repair (NER) by mediating acetylation of ERCC4/XPF, thereby promoting formation of the ERCC4-ERCC1 complex (PubMed:[32034146](#)). Relieves NR1D2-mediated inhibition of APOC3 expression by acetylating NR1D2 (PubMed:[17996965](#)). Acts as a regulator of regulatory T-cells (Treg) by catalyzing FOXP3 acetylation, thereby promoting FOXP3 transcriptional repressor activity (PubMed:[17360565](#), PubMed:[24835996](#)). Involved in skeletal myoblast differentiation by mediating acetylation of SOX4 (PubMed:[26291311](#)). Catalyzes acetylation of APBB1/FE65, increasing its transcription activator activity (PubMed:[33938178](#)). Promotes transcription elongation during the activation phase of the circadian cycle by catalyzing acetylation of BMAL1, promoting elongation of circadian transcripts (By similarity). Together with GSK3 (GSK3A or GSK3B), acts as a regulator of autophagy: phosphorylated at Ser-86 by GSK3 under starvation conditions, leading to activate acetyltransferase activity and promote acetylation of key autophagy regulators, such as ULK1 and RUBCNL/Pacer (PubMed:[30704899](#)). Acts as a regulator of the cGAS-STING innate antiviral response by catalyzing acetylation the N-terminus of CGAS, thereby promoting CGAS DNA-binding and activation (PubMed:[32817552](#)). Also regulates lipid metabolism by mediating acetylation of CHKA or LPIN1 (PubMed:[34077757](#)). Promotes lipolysis of lipid droplets following glucose deprivation by mediating acetylation of isoform 1 of CHKA, thereby promoting monomerization of CHKA and its conversion into a tyrosine-protein kinase (PubMed:[34077757](#)). Acts as a regulator of fatty-acid-induced triacylglycerol synthesis by catalyzing acetylation of LPIN1, thereby promoting the synthesis of diacylglycerol (PubMed:[29765047](#)). In addition to protein acetyltransferase, can use different acyl-CoA substrates, such as (2E)-butenoyl-CoA (crotonyl-CoA), S-lactoyl-CoA (lactyl-CoA) and 2-hydroxyisobutanoyl-CoA (2-hydroxyisobutyryl-CoA), and is able to mediate protein crotonylation, lactylation and 2-hydroxyisobutyrylation, respectively (PubMed:[29192674](#), PubMed:[34608293](#), PubMed:[38961290](#)). Acts as a key regulator of chromosome segregation and kinetochore-microtubule attachment during mitosis by mediating acetylation or crotonylation of

target proteins (PubMed:[26829474](#), PubMed:[29040603](#), PubMed:[30409912](#), PubMed:[34608293](#)). Catalyzes acetylation of AURKB at kinetochores, increasing AURKB activity and promoting accurate chromosome segregation in mitosis (PubMed:[26829474](#)). Acetylates RAN during mitosis, promoting microtubule assembly at mitotic chromosomes (PubMed:[29040603](#)). Acetylates NDC80/HEC1 during mitosis, promoting robust kinetochore-microtubule attachment (PubMed:[30409912](#)). Catalyzes crotonylation of MAPRE1/EB1, thereby ensuring accurate spindle positioning in mitosis (PubMed:[34608293](#)). Catalyzes lactylation of NBN/NBS1 in response to DNA damage, thereby promoting DNA double-strand breaks (DSBs) via homologous recombination (HR) (PubMed:[38961290](#)).

#### Cellular Location


Nucleus. Chromosome. Cytoplasm Chromosome, centromere, kinetochore Cytoplasm, cytoskeleton, spindle pole Nucleus, nucleolus. Cytoplasm, perinuclear region. Note=Upon stimulation with EDN1, it is exported from the nucleus to the perinuclear region and UV irradiation induces translocation into punctuate subnuclear structures named nuclear bodies (PubMed:11262386). Transiently localizes to kinetochores in early mitosis (PubMed:26829474). Localizes to spindle poles when chromosomes align during metaphase (PubMed:34608293). Localizes in the cytoplasm and nucleus of round spermatids (By similarity).  
{ECO:0000250|UniProtKB:Q8CHK4, ECO:0000269|PubMed:11262386, ECO:0000269|PubMed:26829474, ECO:0000269|PubMed:34608293}

#### KAT5 / Tip60/HTATIP Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### KAT5 / Tip60/HTATIP Antibody (C-term) - Images



Western blot analysis of anti-HTATIP Pab (Cat. #AP1081b) in CEM cell line tissue lysate (35ug/lane).HTATIP(arrow) was detected using the purified Pab.

**KAT5 / Tip60/HTATIP Antibody (C-term) - Background**

HTATIP belongs to the MYST family of histone acetyl transferases (HATs) and was originally isolated as an HIV-1 TAT-interactive protein. HATs play important roles in regulating chromatin remodeling, transcription and other nuclear processes by acetylating histone and nonhistone proteins. This protein is a histone acetylase that has a role in DNA repair and apoptosis and is thought to play an important role in signal transduction.

**KAT5 / Tip60/HTATIP Antibody (C-term) - References**

Cai, Y., et al., J. Biol. Chem. 280(14):13665-13670 (2005).  
Patel, J.H., et al., Mol. Cell. Biol. 24(24):10826-10834 (2004).  
Legube, G., et al., J. Biol. Chem. 279(43):44825-44833 (2004).  
Franzese, O., et al., Cell Death Differ. 11(7):782-784 (2004).  
Taubert, S., et al., Mol. Cell. Biol. 24(10):4546-4556 (2004).