

DNAJC22 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10871B

Specification

DNAJC22 Antibody (C-term) - Product Information

Application IHC-P, WB,E Primary Accession Q8N4W6

Other Accession <u>Q170W0</u>, <u>NP 079178.2</u>

Reactivity
Predicted
Host
Clonality
Isotype
Calculated MW
Antigen Region

Human
Bovine
Rabbit
Polyclonal
Rabbit IgG
247-276

DNAJC22 Antibody (C-term) - Additional Information

Gene ID 79962

Other Names

DnaJ homolog subfamily C member 22, DNAJC22

Target/Specificity

This DNAJC22 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 247-276 amino acids from the C-terminal region of human DNAJC22.

Dilution

IHC-P~~1:50~100 WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DNAJC22 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DNAJC22 Antibody (C-term) - Protein Information

Name DNAJC22



Function May function as a co-chaperone.

Cellular Location


Membrane; Multi-pass membrane protein

DNAJC22 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

DNAJC22 Antibody (C-term) - Images



DNAJC22 Antibody (C-term) (Cat. #AP10871b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the DNAJC22 antibody detected the DNAJC22 protein (arrow).



DNAJC22 antibody (C-term) (Cat. #AP10871b) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the DNAJC22 antibody (C-term) for





immunohistochemistry. Clinical relevance has not been evaluated.

DNAJC22 Antibody (C-term) - Background

May function as a co-chaperone.

DNAJC22 Antibody (C-term) - References

Behr, M., et al. Nat. Cell Biol. 9(7):847-853(2007)