

IQCI Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10910b

Specification

IQCJ Antibody (C-term) - Product Information

Application FC, IHC-P, WB,E

Primary Accession 01A5X6

Other Accession NP 001036170.1 Human, Mouse Reactivity

Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 18226

Antigen Region 98-126

IQCJ Antibody (C-term) - Additional Information

Gene ID 654502

Other Names

IQ domain-containing protein J, IQCJ

Target/Specificity

This IQCI antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 98-126 amino acids from the C-terminal region of human IQCJ.

Dilution

FC~~1:10~50 IHC-P~~1:50~100 WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IQC| Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

IQCJ Antibody (C-term) - Protein Information

Name IQCJ (HGNC:32406)




IQCJ Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

IQCJ Antibody (C-term) - Images




IQCJ Antibody (C-term) (Cat. #AP10910b) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the IQCJ antibody detected the IQCJ protein (arrow).




IQCJ Antibody (C-term) (Cat. #AP10910b) western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane). This demonstrates the IQCJ antibody detected the IQCJ protein (arrow).





IQCJ Antibody (C-term) (Cat. #AP10910b) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the IQCJ Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



IQCJ Antibody (C-term) (Cat. #AP10910b) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

IQCJ Antibody (C-term) - References

Martin, P.M., et al. J. Neurosci. 28(24):6111-6117(2008) Kwasnicka-Crawford, D.A., et al. Biochem. Biophys. Res. Commun. 350(4):890-899(2006)