

ZNF674 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10912a**Specification**

ZNF674 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q2M3X9
Other Accession	Q5JW0 , NP_001034980.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	67199
Antigen Region	74-102

ZNF674 Antibody (N-term) - Additional Information**Gene ID** 641339**Other Names**

Zinc finger protein 674, ZNF674

Target/Specificity

This ZNF674 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 74-102 amino acids from the N-terminal region of human ZNF674.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZNF674 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF674 Antibody (N-term) - Protein Information**Name** ZNF674**Function** May be involved in transcriptional regulation.

Cellular Location

Nucleus.


Tissue Location

Expressed in testis..

ZNF674 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF674 Antibody (N-term) - Images

ZNF674 Antibody (N-term) (Cat. #AP10912a) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the ZNF674 antibody detected the ZNF674 protein (arrow).

ZNF674 Antibody (N-term) - Background

This gene encodes a zinc finger protein with an N-terminal Kruppel-associated box-containing (KRAB) domain and 11 Kruppel-type C2H2 zinc finger domains. Like other zinc finger proteins, this gene may function as a transcription factor. This gene resides on an area of chromosome X that has been implicated in nonsyndromic X-linked mental retardation. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq].

ZNF674 Antibody (N-term) - References

Lugtenberg, D., et al. Am. J. Hum. Genet. 78(2):265-278(2006)
Venter, J.C., et al. Science 291(5507):1304-1351(2001)