

UPP2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10995b

Specification

UPP2 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O95045
Other Accession	Q8CGR7 , NP_001128570.1 , NP_775491.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35527
Antigen Region	217-244

UPP2 Antibody (C-term) - Additional Information

Gene ID 151531

Other Names

Uridine phosphorylase 2, UPase 2, UrdPase 2, UPP2

Target/Specificity

This UPP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 217-244 amino acids from the C-terminal region of human UPP2.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UPP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UPP2 Antibody (C-term) - Protein Information

Name UPP2 ([HGNC:23061](#))

Function Catalyzes the reversible phosphorylytic cleavage of uridine to uracil and

ribose-1-phosphate which can then be utilized as carbon and energy sources or in the rescue of pyrimidine bases for nucleotide synthesis (PubMed:[12849978](#), PubMed:[21855639](#)). Shows broad substrate specificity and can also accept deoxyuridine and other analogous compounds (PubMed:[12849978](#)).

Tissue Location


Predominantly expressed in kidney.

UPP2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UPP2 Antibody (C-term) - Images



UPP2 Antibody (C-term) (Cat. #AP10995b) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the UPP2 antibody detected the UPP2 protein (arrow).



UPP2 antibody (C-term) (Cat. #AP10995b) immunohistochemistry analysis in formalin fixed and

paraffin embedded human kidney carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the UPP2 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

UPP2 Antibody (C-term) - Background

Catalyzes the reversible phosphorylytic cleavage of uridine and deoxyuridine to uracil and ribose-or deoxyribose-1-phosphate. The produced molecules are then utilized as carbon and energy sources or in the rescue of pyrimidine bases for nucleotide synthesis. Shows substrate specificity and accept uridine, deoxyuridine, and thymidine as well as the two pyrimidine nucleoside analogs 5-fluorouridine and 5-fluoro-2(')-deoxyuridine as substrates.

UPP2 Antibody (C-term) - References

Maestrini, E., et al. Mol. Psychiatry 15(9):954-968(2010)
Lamesch, P., et al. Genomics 89(3):307-315(2007)
Johansson, M. Biochem. Biophys. Res. Commun. 307(1):41-46(2003)
Russell, R.L., et al. J. Biol. Chem. 276(16):13302-13307(2001)