

TIGD3 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP11042b

Specification

TIGD3 Antibody (C-term) - Product Information

Application WB, IHC-P, FC,E **Primary Accession 06B0B8** Other Accession NP 663771.1 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 52027 Antigen Region 378-407

TIGD3 Antibody (C-term) - Additional Information

Gene ID 220359

Other Names

Tigger transposable element-derived protein 3, TIGD3

Target/Specificity

This TIGD3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 378-407 amino acids from the C-terminal region of human TIGD3.

Dilution

WB~~1:1000 IHC-P~~1:50~100 FC~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TIGD3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TIGD3 Antibody (C-term) - Protein Information

Name TIGD3



Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00320, ECO:0000255|PROSITE-ProRule:PRU00583}

TIGD3 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

TIGD3 Antibody (C-term) - Images




TIGD3 Antibody (C-term) (Cat. #AP11042b) western blot analysis in HepG2 and HL-60 cell line lysates (35ug/lane). This demonstrates the TIGD3 antibody detected the TIGD3 protein (arrow).



TIGD3 antibody (C-term) (Cat. #AP11042b) immunohistochemistry analysis in formalin fixed and paraffin embedded human hodgkin lymphoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TIGD3 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.





TIGD3 Antibody (C-term) (Cat. #AP11042b) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

TIGD3 Antibody (C-term) - Background

The protein encoded by this gene belongs to the tigger subfamily of the pogo superfamily of DNA-mediated transposons in humans. These proteins are related to DNA transposons found in fungi and nematodes, and more distantly to the Tc1 and mariner transposases. They are also very similar to the major mammalian centromere protein B. The exact function of this gene is not known.

TIGD3 Antibody (C-term) - References

Gerhard, D.S., et al. Genome Res. 14 (10B), 2121-2127 (2004) : Bonaldo, M.F., et al. Genome Res. 6(9):791-806(1996)