

UBFD1 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11074a**Specification**

UBFD1 Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	O14562
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	33382
Antigen Region	1-30

UBFD1 Antibody (N-term) - Additional Information**Gene ID** 56061**Other Names**

Ubiquitin domain-containing protein UBFD1, Ubiquitin-binding protein homolog, UBFD1, UBPH

Target/Specificity

This UBFD1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human UBFD1.

Dilution

WB~~1:1000

IHC-P~~1:50~100

FC~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UBFD1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UBFD1 Antibody (N-term) - Protein Information**Name** UBFD1

Synonyms UBPH


Function May play a role as NF-kappa-B regulator.

UBFD1 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UBFD1 Antibody (N-term) - Images




UBFD1 Antibody (N-term) (Cat. #AP11074a) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the UBFD1 antibody detected the UBFD1 protein (arrow).



UBFD1 antibody (N-term) (Cat. #AP11074a) immunohistochemistry analysis in formalin fixed and paraffin embedded human bladder carcinoma followed by peroxidase conjugation of the

secondary antibody and DAB staining. This data demonstrates the use of UBFD1 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



UBFD1 Antibody (N-term) (Cat. #AP11074a) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.