

ELP3 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1112b**Specification**

ELP3 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9H9T3
Other Accession	O9CZX0 , O2KJ61
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	62259
Antigen Region	468-498

ELP3 Antibody (C-term) - Additional Information**Gene ID** 55140**Other Names**

Elongator complex protein 3, hELP3, ELP3

Target/Specificity

This ELP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 468-498 amino acids from the C-terminal region of human ELP3.

Dilution

WB~~1:1000

IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ELP3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ELP3 Antibody (C-term) - Protein Information**Name** ELP3 {ECO:0000303|PubMed:15902492, ECO:0000312|HGNC:HGNC:20696}

Function Catalytic tRNA acetyltransferase subunit of the elongator complex which is required for multiple tRNA modifications, including mcm5U (5-methoxycarbonylmethyl uridine), mcm5s2U (5-methoxycarbonylmethyl-2-thiouridine), and ncm5U (5-carbamoylmethyl uridine) (PubMed:[29415125](#)). In the elongator complex, acts as a tRNA uridine(34) acetyltransferase by mediating formation of carboxymethyluridine in the wobble base at position 34 in tRNAs (By similarity). May also act as a protein lysine acetyltransferase by mediating acetylation of target proteins; such activity is however unclear in vivo and recent evidences suggest that ELP3 primarily acts as a tRNA acetyltransferase (PubMed:[29415125](#)). Involved in neurogenesis: regulates the migration and branching of projection neurons in the developing cerebral cortex, through a process depending on alpha-tubulin acetylation (PubMed:[19185337](#)). Required for acetylation of GJA1 in the developing cerebral cortex (By similarity).

Cellular Location

Cytoplasm. Nucleus [Isoform 2]: Cytoplasm. Nucleus

Tissue Location


Expressed in the cerebellum and spinal motor neurons.

ELP3 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ELP3 Antibody (C-term) - Images



Western blot analysis of ELP3 (arrow) using rabbit polyclonal ELP3 Antibody (C-term) (Cat.#AP1112b).293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the ELP3 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human testis tissue reacted with ELP3 antibody (C-term) (Cat.#AP1112b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

ELP3 Antibody (C-term) - Background

Elongator complex protein 3 (ELP3) is a catalytic histone acetyltransferase subunit of the RNA polymerase II elongator complex, which is a component of the RNA polymerase II (Pol II) holoenzyme and is involved in transcriptional elongation. Elongator may play a role in chromatin remodeling and is involved in acetylation of histones H3 and probably H4. It may also have a methyltransferase activity.

ELP3 Antibody (C-term) - References

Han,Q., Acta Biochim. Biophys. Sin. (Shanghai) 39 (6), 453-461 (2007)
Kim,J.H., Proc. Natl. Acad. Sci. U.S.A. 99 (3), 1241-1246 (2002)
Ninomiya,Y., J. Biochem. 118 (2), 380-389 (1995)