

GOG8A Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11196A

Specification

GOG8A Antibody (N-term) - Product Information

Application	IF, IHC-P, WB,E
Primary Accession	A7E2F4
Other Accession	A8MQT2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	70117
Antigen Region	129-157

GOG8A Antibody (N-term) - Additional Information

Gene ID 23015

Other Names

Golgin subfamily A member 8A, 88 kDa Golgi matrix protein, GM88, GM88 autoantigen, GOLGA8A

Target/Specificity

This GOG8A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 129-157 amino acids from the N-terminal region of human GOG8A.

Dilution

IF~~1:10~50

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GOG8A Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GOG8A Antibody (N-term) - Protein Information

Name GOLGA8A

Synonyms KIAA0855

Function May be involved in maintaining Golgi structure.

Cellular Location


Golgi apparatus, Golgi stack membrane; Peripheral membrane protein

GOG8A Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


GOG8A Antibody (N-term) - Images



Confocal immunofluorescent analysis of GOG8A Antibody (N-term)(Cat#AP11196a) with 293 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



GOG8A Antibody (N-term) (Cat. #AP11196a) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the GOG8A antibody detected the GOG8A protein (arrow).



GOG8A Antibody (N-term) (Cat. #AP11196a) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of GOG8A Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.