

GFER Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11198c**Specification**

GFER Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P55789
Other Accession	NP_005253
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	67-94

GFER Antibody (Center) - Additional Information**Gene ID** 2671**Other Names**

FAD-linked sulfhydryl oxidase ALR, Augmenter of liver regeneration, hERV1, Hepatopoietin, GFER, ALR, HERV1, HPO

Target/Specificity

This GFER antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 67-94 amino acids from the Central region of human GFER.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GFER Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

GFER Antibody (Center) - Protein Information**Name** GFER**Synonyms** ALR, HERV1, HPO

Function [Isoform 1]: FAD-dependent sulfhydryl oxidase that regenerates the redox-active disulfide bonds in CHCHD4/MIA40, a chaperone essential for disulfide bond formation and protein folding in the mitochondrial intermembrane space. The reduced form of CHCHD4/MIA40 forms a transient intermolecular disulfide bridge with GFER/ERV1, resulting in regeneration of the essential disulfide bonds in CHCHD4/MIA40, while GFER/ERV1 becomes re-oxidized by donating electrons to cytochrome c or molecular oxygen.

Cellular Location

[Isoform 1]: Mitochondrion intermembrane space. Mitochondrion

Tissue Location


Ubiquitously expressed. Highest expression in the testis and liver and low expression in the muscle

GFER Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GFER Antibody (Center) - Images



GFER Antibody (Center) (Cat. #AP11198c) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the GFER antibody detected the GFER protein (arrow).



GFER Antibody (Center) (Cat.# AP11198c) western blot analysis in MCF-7,U251 cell line and rat testis tissue lysates (35ug/lane).This demonstrates the GFER antibody detected the GFER protein (arrow).

GFER Antibody (Center) - Background

The hepatotrophic factor designated augments of liver regeneration (ALR) is thought to be one of the factors responsible for the extraordinary regenerative capacity of mammalian liver. It has also been called hepatic regenerative stimulation substance (HSS). The gene resides on chromosome 16 in the interval containing the locus for polycystic kidney disease (PKD1). The putative gene product is 42% similar to the scERV1 protein of yeast. The yeast scERV1 gene had been found to be essential for oxidative phosphorylation, the maintenance of mitochondrial genomes, and the cell division cycle. The human gene is both the structural and functional homolog of the yeast scERV1 gene.

GFER Antibody (Center) - References

Dong, L.Y., et al. Biochem. J. 431(2):277-287(2010) Li, W., et al. FEBS Lett. 584(18):3929-3935(2010) Daithankar, V.N., et al. Biochemistry 49(31):6737-6745(2010) Dayoub, R., et al. Biochem. Biophys. Res. Commun. 395(4):465-470(2010) Chang, J., et al. World J. Gastroenterol. 16(2):193-200(2010)