

**CCNH Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP11210a****Specification**

---

**CCNH Antibody (N-term) - Product Information**

Application	FC, WB,E
Primary Accession	<a href="#">P51946</a>
Other Accession	<a href="#">Q61458</a> , <a href="#">Q4R7U4</a> , <a href="#">Q3ZBL9</a> , <a href="#">NP_001230</a>
Reactivity	Mouse
Predicted	Bovine, Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	37643
Antigen Region	24-52

**CCNH Antibody (N-term) - Additional Information****Gene ID** 902**Other Names**

Cyclin-H, MO15-associated protein, p34, p37, CCNH

**Target/Specificity**

This CCNH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 24-52 amino acids from the N-terminal region of human CCNH.

**Dilution**

FC~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CCNH Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CCNH Antibody (N-term) - Protein Information****Name** CCNH

**Function** Regulates CDK7, the catalytic subunit of the CDK-activating kinase (CAK) enzymatic complex. CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and CDK6 by threonine phosphorylation. CAK complexed to the core-TFIIF basal transcription factor activates RNA polymerase II by serine phosphorylation of the repetitive C-terminal domain (CTD) of its large subunit (POLR2A), allowing its escape from the promoter and elongation of the transcripts. Involved in cell cycle control and in RNA transcription by RNA polymerase II. Its expression and activity are constant throughout the cell cycle.

#### **Cellular Location**


Nucleus.

#### **CCNH Antibody (N-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **CCNH Antibody (N-term) - Images**



CCNH Antibody (N-term) (Cat. #AP11210a) western blot analysis in mouse kidney tissue lysates (35ug/lane). This demonstrates the CCNH antibody detected the CCNH protein (arrow).



CCNH Antibody (N-term) (Cat. #AP11210a) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### **CCNH Antibody (N-term) - Background**

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK7 kinase and ring finger protein MAT1. The kinase complex is able to phosphorylate CDK2 and CDC2 kinases, thus functions as a CDK-activating kinase (CAK). This cyclin and its kinase partner are components of TFIIH, as well as RNA polymerase II protein complexes. They participate in two different transcriptional regulation processes, suggesting an important link between basal transcription control and the cell cycle machinery. A pseudogene of this gene is found on chromosome 4. Alternate splicing results in multiple transcript variants.

### **CCNH Antibody (N-term) - References**

Guey, L.T., et al. Eur. Urol. 57(2):283-292(2010)  
Hosgood, H.D. III, et al. Respir Med 103(12):1866-1870(2009)  
Young, R.P., et al. Postgrad Med J 85(1008):515-524(2009)  
Kweekel, D.M., et al. Br. J. Cancer 101(2):357-362(2009)  
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)