

SERTAD4 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11379b

Specification

SERTAD4 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9NUC0
Other Accession	NP_062551.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	39348
Antigen Region	265-293

SERTAD4 Antibody (C-term) - Additional Information

Gene ID 56256

Other Names

SERTA domain-containing protein 4, SERTAD4

Target/Specificity

This SERTAD4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 265-293 amino acids from the C-terminal region of human SERTAD4.

Dilution

WB~~1:1000

IHC-P~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SERTAD4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SERTAD4 Antibody (C-term) - Protein Information


Name SERTAD4

SERTAD4 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SERTAD4 Antibody (C-term) - Images



Anti-SERTAD4 Antibody (C-term) at 1:1000 dilution + HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



SERTAD4 Antibody (C-term) (Cat. #AP11379b) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of SERTAD4 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.