

TSGA10IP Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11407b

Specification

TSGA10IP Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q3SY00
Other Accession	NP_689975.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	62381
Antigen Region	492-520

TSGA10IP Antibody (C-term) - Additional Information

Gene ID 254187

Other Names

Testis-specific protein 10-interacting protein, Tsga10-interacting protein, TSGA10IP

Target/Specificity

This TSGA10IP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 492-520 amino acids from the C-terminal region of human TSGA10IP.

Dilution

WB~~1:1000

IHC-P~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TSGA10IP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TSGA10IP Antibody (C-term) - Protein Information


Name TSGA10IP

TSGA10IP Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TSGA10IP Antibody (C-term) - Images



TSGA10IP Antibody (C-term) (Cat. #AP11407b) western blot analysis in Ramos cell line lysates (35ug/lane). This demonstrates the TSGA10IP antibody detected the TSGA10IP protein (arrow).



TSGA10IP Antibody (C-term) (Cat. #AP11407b) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TSGA10IP Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

TSGA10IP Antibody (C-term) - References

Strausberg, R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903(2002)