

## **TSGA10IP** Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP11407b

### **Specification**

## TSGA10IP Antibody (C-term) - Product Information

**Application** WB, IHC-P,E **Primary Accession** O3SY00 Other Accession NP 689975.2 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 62381 Antigen Region 492-520

### TSGA10IP Antibody (C-term) - Additional Information

#### Gene ID 254187

#### **Other Names**

Testis-specific protein 10-interacting protein, Tsga10-interacting protein, TSGA10IP

#### Target/Specificity

This TSGA10IP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 492-520 amino acids from the C-terminal region of human TSGA10IP.

# **Dilution**

WB~~1:1000 IHC-P~~1:10~50

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

TSGA10IP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### TSGA10IP Antibody (C-term) - Protein Information

Name TSGA10IP




# TSGA10IP Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# TSGA10IP Antibody (C-term) - Images



TSGA10IP Antibody (C-term) (Cat. #AP11407b) western blot analysis in Ramos cell line lysates (35ug/lane). This demonstrates the TSGA10IP antibody detected the TSGA10IP protein (arrow).



TSGA10IP Antibody (C-term) (Cat. #AP11407b)immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TSGA10IP Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

### TSGA10IP Antibody (C-term) - References





Strausberg, R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903(2002)