

**NT5DC4 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP11408c**

**Specification**

**NT5DC4 Antibody (Center) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q86YG4</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	49027
Antigen Region	172-200

**NT5DC4 Antibody (Center) - Additional Information**

**Gene ID** 284958

**Other Names**

5'-nucleotidase domain-containing protein 4, NT5DC4

**Target/Specificity**

This NT5DC4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 172-200 amino acids from the Central region of human NT5DC4.

**Dilution**

WB~~1:1000

IHC-P~~1:10~50

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NT5DC4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**NT5DC4 Antibody (Center) - Protein Information**


**Name** NT5DC4

## NT5DC4 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## NT5DC4 Antibody (Center) - Images



NT5DC4 Antibody (Center) (Cat. #AP11408c) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the NT5DC4 antibody detected the NT5DC4 protein (arrow).



NT5DC4 Antibody (Center) (Cat. #AP11408c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of NT5DC4 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.