

**KSR2 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP11443b****Specification**

---

**KSR2 Antibody (C-term) - Product Information**

Application	IHC-P, WB, FC,E
Primary Accession	<a href="#">Q6VAB6</a>
Other Accession	<a href="#">NP_775869.3</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	107632
Antigen Region	603-638

**KSR2 Antibody (C-term) - Additional Information****Gene ID** 283455**Other Names**

Kinase suppressor of Ras 2, hKSR2, KSR2

**Target/Specificity**

This KSR2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 603-638 amino acids from the C-terminal region of human KSR2.

**Dilution**

IHC-P~~1:50~100

WB~~1:1000

FC~~1:10~50

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

KSR2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**KSR2 Antibody (C-term) - Protein Information****Name** KSR2 ([HGNC:18610](#))

**Function** Location-regulated scaffold connecting MEK to RAF. Has very low protein kinase activity and can phosphorylate MAP2K1 at several Ser and Thr residues with very low efficiency (in vitro). Acts as MAP2K1/MEK1-dependent allosteric activator of BRAF; upon binding to MAP2K1/MEK1, dimerizes with BRAF and promotes BRAF-mediated phosphorylation of MAP2K1/MEK1 (PubMed:[29433126](#)). Interaction with BRAF enhances KSR2-mediated phosphorylation of MAP2K1 (in vitro). Blocks MAP3K8 kinase activity and MAP3K8-mediated signaling. Acts as a negative regulator of MAP3K3-mediated activation of ERK, JNK and NF- kappa-B pathways, inhibiting MAP3K3-mediated interleukin-8 production.

**Cellular Location**

Cytoplasm. Membrane; Peripheral membrane protein


**Tissue Location**

Mainly expressed in brain and kidney.


**KSR2 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**KSR2 Antibody (C-term) - Images**

KSR2 Antibody (C-term) (Cat. #AP11443b) western blot analysis in 293, K562, HepG2, HeLa cell line lysates (35ug/lane). This demonstrates the KSR2 antibody detected the KSR2 protein (arrow).



KSR2 Antibody (C-term) (Cat. #AP11443b) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of KSR2 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



KSR2 Antibody (C-term) (Cat. #AP11443b) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### **KSR2 Antibody (C-term) - Background**

Location-regulated scaffold connecting MEK to RAF. Blocks MAP3K8 kinase activity and MAP3K8-mediated signaling. Acts as a negative regulator of MAP3K3-mediated activation of ERK, JNK and NF-kappa-B pathways, inhibiting MAP3K3-mediated interleukin-8 production.

### **KSR2 Antibody (C-term) - References**

- Bailey, S.D., et al. Diabetes Care (2010) In press :
- Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
- Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
- Liu, L., et al. Biochim. Biophys. Acta 1794(10):1485-1495(2009)
- Zemunik, T., et al. Croat. Med. J. 50(1):23-33(2009)