

CCDC81 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11477c**Specification**

CCDC81 Antibody (Center) - Product Information

Application	FC, WB,E
Primary Accession	O6ZN84
Other Accession	O9D5W4 , NP_001149946.1 , NP_068599.3
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	76084
Antigen Region	380-408

CCDC81 Antibody (Center) - Additional Information**Gene ID** 60494**Other Names**

Coiled-coil domain-containing protein 81, CCDC81

Target/Specificity

This CCDC81 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 380-408 amino acids from the Central region of human CCDC81.

Dilution

FC~~1:10~50

WB~~1:500

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CCDC81 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CCDC81 Antibody (Center) - Protein Information**Name** CCDC81


Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome


CCDC81 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCDC81 Antibody (Center) - Images

All lanes : Anti-CCDC81 Antibody (Center) at 1:500 dilution Lane 1: HeLa whole cell lysate Lane 2: MDA-MB-453 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 76 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



CCDC81 Antibody (Center) (Cat. #AP11477c) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.