

LMTK3 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11540a

Specification

LMTK3 Antibody (N-term) - Product Information

Application	WB, FC,E
Primary Accession	O96Q04
Other Accession	O5XJV6 , XP_055866.6
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	156-187

LMTK3 Antibody (N-term) - Additional Information

Gene ID 114783

Other Names

Serine/threonine-protein kinase LMTK3, Lemur tyrosine kinase 3, LMTK3, KIAA1883, TYKLM3

Target/Specificity

This LMTK3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 156-187 amino acids from the N-terminal region of human LMTK3.

Dilution

WB~~1:1000-1;2000

FC~~1:25

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LMTK3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

LMTK3 Antibody (N-term) - Protein Information

Name LMTK3

Synonyms KIAA1883, TYKLM3

Function Protein kinase which phosphorylates ESR1 (in vitro) and protects it against proteasomal degradation. May also regulate ESR1 levels indirectly via a PKC-AKT-FOXO3 pathway where it decreases the activity of PKC and the phosphorylation of AKT, thereby increasing binding of transcriptional activator FOXO3 to the ESR1 promoter and increasing ESR1 transcription (PubMed:[21602804](#)). Involved in endocytic trafficking of N-methyl-D-aspartate receptors (NMDAR) in neurons (By similarity).

Cellular Location


Membrane {ECO:0000250|UniProtKB:Q5XJV6}; Single- pass membrane protein {ECO:0000250|UniProtKB:Q5XJV6}. Cell projection, axon {ECO:0000250|UniProtKB:Q5XJV6}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q5XJV6}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q5XJV6}. Note=Punctate pattern in cell projections

LMTK3 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


LMTK3 Antibody (N-term) - Images



LMTK3 Antibody (N-term) (Cat. #AP11540a) western blot analysis in mouse liver tissue lysates (35ug/lane). This demonstrates the LMTK3 antibody detected the LMTK3 protein (arrow).



All lanes : Anti-LMTK3 Antibody (N-term) at 1:1000-1;2000 dilution Lane 1: HeLa lysate Lane 2: mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 154 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing U-2 OS cells stained with AP11540a(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP11540a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OE188374) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

LMTK3 Antibody (N-term) - Background

The specific function of the protein remains unknown.