

OCT3/4 Antibody (C279)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11546a**Specification**

OCT3/4 Antibody (C279) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q01860
Other Accession	Q06416 , NP_002692.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	38571
Antigen Region	265-297

OCT3/4 Antibody (C279) - Additional Information**Gene ID** 5460**Other Names**

POU domain, class 5, transcription factor 1, Octamer-binding protein 3, Oct-3, Octamer-binding protein 4, Oct-4, Octamer-binding transcription factor 3, OTF-3, POU5F1, OCT3, OCT4, OTF3

Target/Specificity

This OCT3/4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 265-297 amino acids from human OCT3/4.

Dilution

WB~~1:1000

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OCT3/4 Antibody (C279) is for research use only and not for use in diagnostic or therapeutic procedures.

OCT3/4 Antibody (C279) - Protein Information**Name** POU5F1

Synonyms OCT3, OCT4, OTF3

Function Transcription factor that binds to the octamer motif (5'- ATTTGCAT-3'). Forms a trimeric complex with SOX2 or SOX15 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early embryogenesis and for embryonic stem cell pluripotency.

Cellular Location

Cytoplasm. Nucleus. Note=Expressed in a diffuse and slightly punctuate pattern. Colocalizes with MAPK8 and MAPK9 in the nucleus. {ECO:0000250|UniProtKB:P20263, ECO:0000269|PubMed:18191611, ECO:0000269|PubMed:19274063, ECO:0000269|PubMed:23024368}


Tissue Location

Expressed in developing brain. Highest levels found in specific cell layers of the cortex, the olfactory bulb, the hippocampus and the cerebellum. Low levels of expression in adult tissues.


OCT3/4 Antibody (C279) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OCT3/4 Antibody (C279) - Images

OCT3/4 Antibody (C279) (Cat. #AP11546a) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the OCT3/4 antibody detected the OCT3/4 protein (arrow).



OCT3/4 Antibody (C279) (Cat. #AP11546a) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of OCT3/4 Antibody (C279) for immunohistochemistry. Clinical relevance has not been evaluated.

OCT3/4 Antibody (C279) - Background

This gene encodes a transcription factor containing a POU homeodomain. This transcription factor plays a role in embryonic development, especially during early embryogenesis, and it is necessary for embryonic stem cell pluripotency. A translocation of this gene with the Ewing's sarcoma gene, t(6;22)(p21;q12), has been linked to tumor formation. Alternative splicing, as well as usage of alternative translation initiation codons, results in multiple isoforms, one of which initiates at a non-AUG (CUG) start codon. Related pseudogenes have been identified on chromosomes 1, 3, 8, 10, and 12.

OCT3/4 Antibody (C279) - References

Abu-Remaileh, M., et al. EMBO J. 29(19):3236-3248(2010)
Ucisik-Akkaya, E., et al. Mol. Hum. Reprod. 16(10):770-777(2010)
Schultz, S.S., et al. Mol. Cell. Biol. 30(18):4521-4534(2010)
Kim, S., et al. Br. J. Cancer 102(2):436-446(2010)
Wang, X., et al. Stem Cells 27(6):1265-1275(2009)