

SCEL Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11564c**Specification**

SCEL Antibody (Center) - Product Information

Application	WB, IHC-P-Leica, FC,E
Primary Accession	O95171
Other Accession	NP_659001.2 , NP_003834.3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	260-289

SCEL Antibody (Center) - Additional Information**Gene ID** 8796**Other Names**

Sciellin, SCEL

Target/Specificity

This SCEL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 260-289 amino acids from the Central region of human SCEL.

Dilution

WB~~1:1000
IHC-P-Leica~~1:500
FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SCEL Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

SCEL Antibody (Center) - Protein Information**Name** SCEL**Function** May function in the assembly or regulation of proteins in the cornified envelope. The LIM

domain may be involved in homotypic or heterotypic associations and may function to localize sciellin to the cornified envelope.

Cellular Location

Cytoplasm. Membrane. Note=May become cross-linked to membrane proteins by transglutaminase

Tissue Location


Highly expressed in esophagus. It is also expressed in keratinocytes, amniotic tissue, foreskin stratum spinosum and stratum granulosum, hair follicle and nail

SCEL Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


SCEL Antibody (Center) - Images




Anti-SCEL Antibody (Center) at 1:1000 dilution + A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 78 kDa Blocking/Dilution buffer: 5% NFDm/TBST.




Anti-SCEL Antibody (Center) at 1:1000 dilution + NCI-H292 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 78 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded human esophagus tissue using AP11564C performed on the Leica® BOND RXm. Samples were incubated with primary antibody(1/500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of paraffin-embedded human skin tissue using AP11564C performed on the Leica® BOND RXm. Samples were incubated with primary antibody(1/500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Overlay histogram showing A431 cells stained with AP11564C(green line). The cells were fixed with 2% paraformaldehyde and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at Room temperature. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

SCEL Antibody (Center) - Background

The protein encoded by this gene is a precursor to the cornified envelope of terminally differentiated keratinocytes. This protein localizes to the periphery of cells and may function in the assembly or regulation of proteins in the cornified envelope. Transcript variants encoding different isoforms exist. A transcript variant utilizing an alternative polyA signal has been described in the literature, but its full-length nature has not been determined.

SCEL Antibody (Center) - References

Pyle, A.L., et al. Cardiovasc. Pathol. 19 (2), E13-E20 (2010) :
Corona, W., et al. Anticancer Res. 24 (3A), 1417-1419 (2004) :

Champliaud, M.F., et al. J. Invest. Dermatol. 121(4):781-785(2003)
Champliaud, M.F., et al. Genomics 70(2):264-268(2000)
Champliaud, M.F., et al. J. Biol. Chem. 273(47):31547-31554(1998)