

#### F11 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP11568c

### **Specification**

## F11 Antibody (Center) - Product Information

**Application** IHC-P, WB,E **Primary Accession** P03951 Other Accession NP 000119.1 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 70109 Antigen Region 281-307

### F11 Antibody (Center) - Additional Information

#### **Gene ID 2160**

#### **Other Names**

Coagulation factor XI, FXI, Plasma thromboplastin antecedent, PTA, Coagulation factor XIa heavy chain, Coagulation factor XIa light chain, F11

#### Target/Specificity

This F11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 281-307 amino acids from the Central region of human F11.

#### **Dilution**

IHC-P~~1:50~100 WB~~1:1000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

F11 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### F11 Antibody (Center) - Protein Information

## Name F11





**Function** Factor XI triggers the middle phase of the intrinsic pathway of blood coagulation by activating factor IX.

Cellular Location Secreted.

#### **Tissue Location**


Isoform 2 is produced by platelets and megakaryocytes but absent from other blood cells

# F11 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

### F11 Antibody (Center) - Images



All lanes : Anti-F11 Antibody (Center) at 1:1000 dilution Lane 1: A549 whole cell lysate Lane 2: Jurkat whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 70 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





F11 Antibody (Center) (Cat. #AP11568c)immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of F11 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

# F11 Antibody (Center) - Background

This gene encodes coagulation factor XI of the blood coagulation cascade. This protein is present in plasma as a zymogen, which is a unique plasma coagulation enzyme because it exists as a homodimer consisting of two identical polypeptide chains linked by disulfide bonds. During activation of the plasma factor XI, an internal peptide bond is cleaved by factor XIIa (or XII) in each of the two chains, resulting in activated factor XIa, a serine protease composed of two heavy and two light chains held together by disulfide bonds. This activated plasma factor XI triggers the middle phase of the intrisic pathway of blood coagulation by activating factor IX. Defects in this factor lead to Rosenthal syndrome, a blood coagulation abnormality. [provided by RefSeq].

## F11 Antibody (Center) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Wong, P.C., et al. Thromb. Haemost. 104(2):302-310(2010) Whelihan, M.F., et al. J. Thromb. Haemost. 8(7):1532-1539(2010) Delluc, A., et al. Thromb. Haemost. 103(6):1161-1169(2010) Barber, M.J., et al. PLoS ONE 5 (3), E9763 (2010):