

**GU CY2D Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP11569c****Specification**

---

**GU CY2D Antibody (Center) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">Q02846</a>
Other Accession	<a href="#">NP_000171.1</a>
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	120059
Antigen Region	540-570

**GU CY2D Antibody (Center) - Additional Information****Gene ID** 3000**Other Names**

Retinal guanylyl cyclase 1, RETGC-1, Guanylate cyclase 2D, retinal, Rod outer segment membrane guanylate cyclase, ROS-GC, GUCY2D, CORD6, GUC1A4, GUC2D, RETGC, RETGC1

**Target/Specificity**

This GUCY2D antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 540-570 amino acids from the Central region of human GUCY2D.

**Dilution**

WB~~1:2000

IHC-P~~1:50~100

FC~~1:25

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GUCY2D Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**GU CY2D Antibody (Center) - Protein Information**

**Name** GUCY2D

**Function** Catalyzes the synthesis of cyclic GMP (cGMP) in rods and cones of photoreceptors. Plays an essential role in phototransduction, by mediating cGMP replenishment (PubMed:[15123990](#), PubMed:[21928830](#), PubMed:[26100624](#), PubMed:[30319355](#), PubMed:[9600905](#)). May also participate in the trafficking of membrane-associated proteins to the photoreceptor outer segment membrane (By similarity).

**Cellular Location**

Photoreceptor outer segment membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein


**Tissue Location**

Retina..


**GUCY2D Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**GUCY2D Antibody (Center) - Images**


GUCY2D Antibody (Center) (Cat. #AP11569c) western blot analysis in Ramos cell line lysates (35ug/lane). This demonstrates the GUCY2D antibody detected the GUCY2D protein (arrow).



Anti-GUCY2D Antibody (Center) at 1:2000 dilution + rat eyeball lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 120 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



GUCY2D Antibody (Center) (Cat. #AP11569c) immunohistochemistry analysis in formalin fixed and paraffin embedded human thyroid tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of GUCY2D Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Overlay histogram showing Hela cells stained with AP11569c (green line). The cells were fixed with 2% paraformaldehyde (10 min). The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP11569c, 1:25 dilution)

for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10, 000 events was performed.

### **GUCY2D Antibody (Center) - Background**

This gene encodes a retina-specific guanylate cyclase, which is a member of the membrane guanylyl cyclase family. Like other membrane guanylyl cyclases, this enzyme has a hydrophobic amino-terminal signal sequence followed by a large extracellular domain, a single membrane spanning domain, a kinase homology domain, and a guanylyl cyclase catalytic domain. In contrast to other membrane guanylyl cyclases, this enzyme is not activated by natriuretic peptides. Mutations in this gene result in Leber congenital amaurosis and cone-rod dystrophy-6 diseases. [provided by RefSeq].

### **GUCY2D Antibody (Center) - References**

Silva, L.K., et al. Eur. J. Hum. Genet. 18(11):1221-1227(2010)  
Liu, C.Y., et al. Carcinogenesis 31(7):1259-1263(2010)  
Joslyn, G., et al. Alcohol. Clin. Exp. Res. 34(5):800-812(2010)  
Pasadhika, S., et al. Invest. Ophthalmol. Vis. Sci. 51(5):2608-2614(2010)  
Sundaresan, P., et al. Mol. Vis. 15, 1781-1787 (2009) :