

**DUX4 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP11636C****Specification**

---

**DUX4 Antibody (Center) - Product Information**

Application	WB, FC,E
Primary Accession	<a href="#">O9UBX2</a>
Other Accession	<a href="#">O6RFH8</a> , <a href="#">POCJ90</a> , <a href="#">POCJ89</a> , <a href="#">POCJ88</a> , <a href="#">POCJ87</a> , <a href="#">POCJ86</a> , <a href="#">POCJ85</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	246-275

**DUX4 Antibody (Center) - Additional Information****Gene ID** 100288687**Other Names**

Double homeobox protein 4 {ECO:0000312|HGNC:HGNC:50800}, Double homeobox protein 10 {ECO:0000312|EMBL:AAK915091}, DUX4 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=50800" target="\_blank">HGNC:50800</a>), DUX10

**Target/Specificity**

This DUX4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 246-275 amino acids from the Central region of human DUX4.

**Dilution**

WB~~1:2000

FC~~1:25

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

DUX4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**DUX4 Antibody (Center) - Protein Information**

**Name** DUX4 ([HGNC:50800](#))

**Synonyms** DUX10

**Function** [Isoform 1]: Transcription factor that is selectively and transiently expressed in cleavage-stage embryos (PubMed:[28459457](#)). Binds to double-stranded DNA elements with the consensus sequence 5'- TAATCTAATCA-3' (PubMed:[28459454](#), PubMed:[28459457](#), PubMed:[29572508](#), PubMed:[30315230](#), PubMed:[30540931](#)). Binds to chromatin containing histone H3 acetylated at 'Lys-27' (H3K27ac) and promotes deacetylation of H3K27ac. In parallel, binds to chromatin that lacks histone H3 acetylation at 'Lys-27' (H3K27ac) and recruits EP300 and CREBBP to promote acetylation of histone H3 at 'Lys-27' at new sites (PubMed:[26951377](#)). Involved in transcriptional regulation of numerous genes, primarily as transcriptional activator, but also mediates repression of a set of target genes (PubMed:[17984056](#), PubMed:[26951377](#), PubMed:[27378237](#), PubMed:[28459454](#), PubMed:[28459457](#), PubMed:[29572508](#), PubMed:[29618456](#), PubMed:[30540931](#)). Promotes expression of ZSCAN4 and KDM4E, two proteins with essential roles during early embryogenesis (PubMed:[26951377](#), PubMed:[27378237](#), PubMed:[28459457](#), PubMed:[29618456](#)). Promotes nuclear translocation of CTNNB1/beta-catenin and its subsequent activation of target genes (PubMed:[36158201](#)). Heterologous expression in cultured embryonic stem cells mediates transcription of HERV retrotransposons and transcripts derived from ACRO1 and HSATII satellite repeats (PubMed:[28459457](#)). May activate expression of PITX1 (PubMed:[17984056](#)). May regulate microRNA (miRNA) expression (PubMed:[24145033](#)). Inappropriate expression can inhibit myogenesis and promote apoptosis (PubMed:[26951377](#), PubMed:[28935672](#), PubMed:[29618456](#)).

#### Cellular Location

[Isoform 1]: Nucleus {ECO:0000255|PROSITE- ProRule:PRU00108, ECO:0000269|PubMed:15709750, ECO:0000269|PubMed:17984056, ECO:0000269|PubMed:21060811, ECO:0000269|PubMed:26816005, ECO:0000269|PubMed:26951377, ECO:0000269|PubMed:27378237, ECO:0000269|PubMed:28459457, ECO:0000269|PubMed:29618456}. Cytoplasm Note=Actively transported through the nuclear pore complex (NPC)

#### Tissue Location


Isoform 1: Does not seem to be expressed in normal muscle, but is detected in muscle of individuals with FSHD, and also in testis (at protein level) (PubMed:17984056, PubMed:21060811). Isoform 1: Does not seem to be expressed in normal muscle, but in muscle of individuals with FSHD, where it may be toxic to cells (PubMed:17984056, PubMed:21060811). Isoform 2: Detected in skeletal muscle, fibroblasts and testis from healthy individuals (PubMed:21060811)


#### DUX4 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### DUX4 Antibody (Center) - Images






DUX4 Antibody (Center) (Cat. #AP11636c) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the DUX4 antibody detected the DUX4 protein (arrow).



Anti-DUX4 Antibody (Center) at 1:2000 dilution + Human testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing U-2 OS cells stained with AP11636c (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP11636c, 1:25 dilution) for 60 min at 37°C. The secondary

antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10, 000 events was performed.

**DUX4 Antibody (Center) - Background**

DUX4 may be involved in transcriptional regulation.