

PCDH20 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11655C

Specification

PCDH20 Antibody (Center) - Product Information

Application	WB, IF, FC, IHC-P,E
Primary Accession	Q8N6Y1
Other Accession	NP_073754.2
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	104919
Antigen Region	445-472

PCDH20 Antibody (Center) - Additional Information

Gene ID 64881

Other Names

Protocadherin-20, Protocadherin-13, PCDH20, PCDH13

Target/Specificity

This PCDH20 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 445-472 amino acids from the Central region of human PCDH20.

Dilution

WB~~1:2000
IF~~1:10~50
FC~~1:10~50
IHC-P~~1:10~50
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PCDH20 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PCDH20 Antibody (Center) - Protein Information

Name PCDH20

Synonyms PCDH13

Function Potential calcium-dependent cell-adhesion protein.

Cellular Location


Cell membrane; Single-pass type I membrane protein

PCDH20 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PCDH20 Antibody (Center) - Images




Confocal immunofluorescent analysis of PCDH20 Antibody (Center)(Cat#AP11655c) with MCF-7 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green).DAPI was used to stain the cell nuclear (blue).




All lanes : Anti-PCDH20 Antibody (Center) at 1:1000 dilution Lane 1: SK-BR-3 whole cell lysate Lane 2: SH-SY5Y whole cell lysate Lane 3: MCF-7 whole cell lysate Lane 4: HL-60 whole cell lysate Lane 5: C6 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 105 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-PCDH20 Antibody (Center) at 1:2000 dilution Lane 1: SK-BR-3 whole cell lysate Lane 2: SH-SY5Y whole cell lysate Lane 3: MCF-7 whole cell lysate Lane 4: HL-60 whole cell lysate Lane 5: Rat brain whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 105 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



PCDH20 Antibody (Center) (Cat. #AP11655c) immunohistochemistry analysis in formalin fixed and paraffin embedded human pancreas tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PCDH20 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



PCDH20 Antibody (Center) (Cat. #AP11655c) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

PCDH20 Antibody (Center) - Background

This gene belongs to the protocadherin gene family, a subfamily of the cadherin superfamily. This gene encodes a protein which contains 6 extracellular cadherin domains, a transmembrane domain and a cytoplasmic tail differing from those of the classical cadherins. Although its specific function is undetermined, the cadherin-related neuronal receptor is thought to play a role in the establishment and function of specific cell-cell connections in the brain.

PCDH20 Antibody (Center) - References

- Suzuki, S.T. Exp. Cell Res. 261(1):13-18(2000)
- Nollet, F., et al. J. Mol. Biol. 299(3):551-572(2000)
- Yagi, T., et al. Genes Dev. 14(10):1169-1180(2000)
- Wu, Q., et al. Proc. Natl. Acad. Sci. U.S.A. 97(7):3124-3129(2000)
- Cross, S.H., et al. Nat. Genet. 6(3):236-244(1994)