

PTPN20A Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP11798c

Specification

PTPN20A Antibody (Center) - Product Information

Application FC, IHC-P, WB,E

Primary Accession Q4IDL3

Other Accession <u>NP_001035817.1</u>, <u>NP_001035853.1</u>,

NP 001035849.1

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Polyclonal
Rabbit IgG
A8423
178-207

PTPN20A Antibody (Center) - Additional Information

Gene ID 26095

Other Names

Tyrosine-protein phosphatase non-receptor type 20, hPTPN20, PTPN20A

Target/Specificity

This PTPN20A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 178-207 amino acids from the Central region of human PTPN20A.

Dilution

FC~~1:10~50 IHC-P~~1:10~50 WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PTPN20A Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PTPN20A Antibody (Center) - Protein Information



Name PTPN20 (HGNC:23423)

Function Tyrosine-protein phosphatase targeted to sites of actin polymerization in response of varied extracellular stimuli. Has tyrosine phosphatase activity towards various tyrosyl phosphorylated substrates.

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Colocalizes with the microtubule-organizing center and intracellular membrane compartments

Tissue Location


Present in many cell lines (at protein level). Widely expressed.

PTPN20A Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

PTPN20A Antibody (Center) - Images




PTPN20A Antibody (Center) (Cat. #AP11798c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the PTPN20A antibody detected the PTPN20A protein (arrow).






PTPN20A Antibody (Center) (Cat. #AP11798c)immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PTPN20A Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



PTPN20A Antibody (Center) (Cat. #AP11798c) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

PTPN20A Antibody (Center) - Background

The product of this gene belongs to the family of classical tyrosine-specific protein tyrosine phosphatases. Many protein tyrosine phosphatases have been shown to regulate fundamental cellular processes and several are mutated in human diseases. Chromosome 10q contains a segmental duplication resulting in multiple copies of the protein tyrosine phosphatase, non-receptor type 20 gene. The two nearly identical copies are designated as PTPN20A and PTPN20B. A third copy is only partially duplicated and contains a pseudogene, designated as PTPN20C. This gene encodes the more telomeric copy, PTPN20B. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.





PTPN20A Antibody (Center) - References

Matsuoka, S., et al. Science 316(5828):1160-1166(2007) Fodero-Tavoletti, M.T., et al. Biochem. J. 389 (PT 2), 343-354 (2005) :