

**PELP1 Antibody (Center R759)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP11872c**

**Specification**

---

**PELP1 Antibody (Center R759) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q8IZL8</a>
Other Accession	<a href="#">Q56B11</a> , <a href="#">Q9DBD5</a> , <a href="#">NP_055204</a>
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	744-771

**PELP1 Antibody (Center R759) - Additional Information**

**Gene ID** 27043

**Other Names**

Proline-, glutamic acid- and leucine-rich protein 1, Modulator of non-genomic activity of estrogen receptor, Transcription factor HMX3, PELP1, HMX3, MNAR

**Target/Specificity**

This PELP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 744-771 amino acids from the Central region of human PELP1.

**Dilution**

WB~~1:1000

IHC-P~~1:25

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PELP1 Antibody (Center R759) is for research use only and not for use in diagnostic or therapeutic procedures.

**PELP1 Antibody (Center R759) - Protein Information**

**Name** PELP1

**Synonyms** HMX3, MNAR

**Function** Coactivator of estrogen receptor-mediated transcription and a corepressor of other nuclear hormone receptors and sequence-specific transcription factors (PubMed:[14963108](#)). Plays a role in estrogen receptor (ER) genomic activity when present in the nuclear compartment by activating the ER target genes in a hormonal stimulation dependent manner. Can facilitate ER non-genomic signaling via SRC and PI3K interaction in the cytosol. Plays a role in E2-mediated cell cycle progression by interacting with RB1. May have important functional implications in ER/growth factor cross-talk. Interacts with several growth factor signaling components including EGFR and HRS. Functions as the key stabilizing component of the Five Friends of Methylated CHTOP (5FMC) complex; the 5FMC complex is recruited to ZNF148 by methylated CHTOP, leading to desumoylation of ZNF148 and subsequent transactivation of ZNF148 target genes. Component of the PELP1 complex involved in the nucleolar steps of 28S rRNA maturation and the subsequent nucleoplasmic transit of the pre-60S ribosomal subunit. Regulates pre-60S association of the critical remodeling factor MDN1 (PubMed:[21326211](#)). May promote tumorigenesis via its interaction with and modulation of several oncogenes including SRC, PI3K, STAT3 and EGFR. Plays a role in cancer cell metastasis via its ability to modulate E2-mediated cytoskeleton changes and cell migration via its interaction with SRC and PI3K.

**Cellular Location**

Nucleus, nucleolus. Nucleus, nucleoplasm. Nucleus. Cytoplasm Note=Mainly found in the nucleoplasm, with low levels detected in the cytoplasm (By similarity). Also found associated with the plasma membrane. Mainly in cytoplasm in a subset of breast tumors Localization is widely deregulated in endometrial cancers with predominantly cytoplasm localization in high-grade endometrial tumors (PubMed:16140940). {ECO:0000250|UniProtKB:Q9DBD5, ECO:0000269|PubMed:16140940}


**Tissue Location**


Widely expressed..

**PELP1 Antibody (Center R759) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**PELP1 Antibody (Center R759) - Images**




PELP1 Antibody (Center R759) (Cat. #AP11872c) western blot analysis in mouse spleen tissue lysates (35ug/lane). This demonstrates the PELP1 antibody detected the PELP1 protein (arrow).



All lanes : Anti-PELP1 Antibody (Center R759) at 1:1000 dilution Lane 1: T47D whole cell lysate Lane 2: mouse testis lysate Lane 3: Hela whole cell lysate Lane 4: MCF-7 whole cell lysate Lane 5: NIH/3T3 whole cell lysate Lane 6: SW620 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 120 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP11872C staining PELP1 in human colon tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



AP11872C staining PELP1 in human colon tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

#### **PELP1 Antibody (Center R759) - Background**

PELP1 is a coactivator of estrogen receptor (see ESR1; MIM 133430)-mediated transcription and a corepressor of other nuclear hormone receptors and sequence-specific transcription factors (Choi et al., 2004 [PubMed 15456770]).

#### **PELP1 Antibody (Center R759) - References**

Nair, S.S., et al. EMBO Rep. 11(6):438-444(2010)  
Habashy, H.O., et al. Breast Cancer Res. Treat. 120(3):603-612(2010)  
Vadlamudi, R.K., et al. J. Steroid Biochem. Mol. Biol. 118 (4-5), 211-218 (2010) :  
Tzelepi, V., et al. Virchows Arch. 454(1):41-53(2009)  
Grivas, P.D., et al. Cell. Oncol. 31(3):235-247(2009)