

STK32A Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11949a**Specification**

STK32A Antibody (N-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	Q8WU08
Other Accession	NP_659438.1
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	46369
Antigen Region	1-30

STK32A Antibody (N-term) - Additional Information**Gene ID** 202374**Other Names**

Serine/threonine-protein kinase 32A, Yet another novel kinase 1, STK32A, YANK1

Target/Specificity

This STK32A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human STK32A.

Dilution

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

STK32A Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

STK32A Antibody (N-term) - Protein Information**Name** STK32A


Synonyms YANK1**Cellular Location**

Cell membrane; Lipid-anchor


STK32A Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STK32A Antibody (N-term) - Images

STK32A Antibody (N-term) (Cat. #AP11949a) western blot analysis in mouse spleen tissue lysates (35ug/lane). This demonstrates the STK32A antibody detected the STK32A protein (arrow).



STK32A Antibody (N-term) (Cat. #AP11949a) immunohistochemistry analysis in formalin fixed and paraffin embedded human bladder carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of STK32A Antibody

(N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

STK32A Antibody (N-term) - Background

STK32A is a member of the serine/threonine protein kinase family. The specific function of this kinase is not known.

STK32A Antibody (N-term) - References

Rose, J. Phd, et al. Mol. Med. (2010) In press :
Koskinen, L.L., et al. Tissue Antigens 74(5):408-416(2009)
Schmutz, J., et al. Nature 431(7006):268-274(2004)
Manning, G., et al. Science 298(5600):1912-1934(2002)